

Yes1 Antibody
Rabbit mAb
Catalog # AP91854

Specification

Yes1 Antibody - Product Information

| | |
|-----------------------------------|------------------------|
| Application | WB, IHC, ICC, IP |
| Primary Accession | P07947 |
| Reactivity | Rat |
| Clonality | Monoclonal |
| Other Names | |
| c Yes; HsT441; P61 YES;Yes; YES1; | |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 60801 Da |

Yes1 Antibody - Additional Information

| | |
|------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human Yes1 |
| Description | Promotes infectivity of <i>Neisseria gonorrhoeae</i> in epithelial cells by phosphorylating MCP/CD46. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Yes1 Antibody - Protein Information

Name YES1

Synonyms YES

Function

Non-receptor protein tyrosine kinase that is involved in the regulation of cell growth and survival, apoptosis, cell-cell adhesion, cytoskeleton remodeling, and differentiation. Stimulation by receptor tyrosine kinases (RTKs) including EGFR, PDGFR, CSF1R and FGFR leads to recruitment of YES1 to the phosphorylated receptor, and activation and phosphorylation of downstream substrates. Upon EGFR activation, promotes the phosphorylation of PARD3 to favor epithelial tight junction assembly. Participates in the phosphorylation of specific junctional components such as CTNND1 by stimulating the FYN and FER tyrosine kinases at cell-cell contacts. Upon T-cell stimulation by CXCL12, phosphorylates collapsin response mediator protein 2/DPYSL2 and induces T-cell migration. Participates in CD95L/FASLG signaling pathway and mediates AKT-mediated cell migration. Plays a role in cell cycle progression by phosphorylating the cyclin-dependent kinase 4/CDK4 thus regulating the G1 phase. Also involved in G2/M progression and cytokinesis.

Catalyzes phosphorylation of organic cation transporter OCT2 which induces its transport activity (PubMed:26979622).

Cellular Location

Cell membrane. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytosol Note=Newly synthesized protein initially accumulates in the Golgi region and traffics to the plasma membrane through the exocytic pathway

Tissue Location


Expressed in the epithelial cells of renal proximal tubules and stomach as well as hematopoietic cells in the bone marrow and spleen in the fetal tissues. In adult, expressed in epithelial cells of the renal proximal tubules and present in keratinocytes in the basal epidermal layer of epidermis.

Yes1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Yes1 Antibody - Images



Western blot analysis of Yes1 expression in SW480 cell lysate.