

**Cytochrome P450 17A1 Antibody**  
Rabbit mAb  
Catalog # AP91937**Specification****Cytochrome P450 17A1 Antibody - Product Information**

Application	WB, FC, ICC, IP
Primary Accession	<a href="#">P05093</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
CPT7; CYP17; P450C17; S17AH; CYP17A1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	57371 Da

**Cytochrome P450 17A1 Antibody - Additional Information**

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Cytochrome P450 17A1
Description	Conversion of pregnenolone and progesterone to their 17-alpha-hydroxylated products and subsequently to dehydroepiandrosterone (DHEA) and androstenedione.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**Cytochrome P450 17A1 Antibody - Protein Information**

**Name** CYP17A1 {ECO:0000303|PubMed:19793597, ECO:0000312|HGNC:HGNC:2593}

**Function**

A cytochrome P450 monooxygenase involved in corticoid and androgen biosynthesis (PubMed:<a href="http://www.uniprot.org/citations/22266943" target="\_blank">22266943</a>, PubMed:<a href="http://www.uniprot.org/citations/25301938" target="\_blank">25301938</a>, PubMed:<a href="http://www.uniprot.org/citations/27339894" target="\_blank">27339894</a>, PubMed:<a href="http://www.uniprot.org/citations/9452426" target="\_blank">9452426</a>). Catalyzes 17-alpha hydroxylation of C21 steroids, which is common for both pathways. A second oxidative step, required only for androgen synthesis, involves an acyl-carbon cleavage. The 17-alpha hydroxy intermediates, as part of adrenal glucocorticoids biosynthesis pathway, are precursors of cortisol (Probable) (PubMed:<a href="http://www.uniprot.org/citations/25301938" target="\_blank">25301938</a>, PubMed:<a href="http://www.uniprot.org/citations/9452426" target="\_blank">9452426</a>). Hydroxylates steroid hormones, pregnenolone and progesterone

to form 17-alpha hydroxy metabolites, followed by the cleavage of the C17-C20 bond to form C19 steroids, dehydroepiandrosterone (DHEA) and androstenedione (PubMed:<a href="http://www.uniprot.org/citations/22266943" target="\_blank">22266943</a>, PubMed:<a href="http://www.uniprot.org/citations/25301938" target="\_blank">25301938</a>, PubMed:<a href="http://www.uniprot.org/citations/27339894" target="\_blank">27339894</a>, PubMed:<a href="http://www.uniprot.org/citations/36640554" target="\_blank">36640554</a>, PubMed:<a href="http://www.uniprot.org/citations/9452426" target="\_blank">9452426</a>). Has 16-alpha hydroxylase activity. Catalyzes 16-alpha hydroxylation of 17-alpha hydroxy pregnenolone, followed by the cleavage of the C17-C20 bond to form 16-alpha-hydroxy DHEA (PubMed:<a href="http://www.uniprot.org/citations/36640554" target="\_blank">36640554</a>). Also 16-alpha hydroxylates androgens, relevant for estriol synthesis (PubMed:<a href="http://www.uniprot.org/citations/25301938" target="\_blank">25301938</a>, PubMed:<a href="http://www.uniprot.org/citations/27339894" target="\_blank">27339894</a>). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (PubMed:<a href="http://www.uniprot.org/citations/22266943" target="\_blank">22266943</a>, PubMed:<a href="http://www.uniprot.org/citations/25301938" target="\_blank">25301938</a>, PubMed:<a href="http://www.uniprot.org/citations/27339894" target="\_blank">27339894</a>, PubMed:<a href="http://www.uniprot.org/citations/9452426" target="\_blank">9452426</a>).

#### Cellular Location

Endoplasmic reticulum membrane. Microsome membrane


#### Cytochrome P450 17A1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Cytochrome P450 17A1 Antibody - Images





Western blot analysis of Cytochrome P450 17A1 expression in Jurkat cell lysate.