

DDB2 Antibody
Rabbit mAb
Catalog # AP92039

Specification

DDB2 Antibody - Product Information

Application	WB, IHC, FC, ICC
Primary Accession	O92466
Clonality	Monoclonal
Other Names	
DDB p48 subunit; Ddb2; DDBb; UV-DDB 2;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	47864 Da

DDB2 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human DDB2
Description	Required for DNA repair. Binds to DDB1 to form the UV-damaged DNA-binding protein complex (the UV-DDB complex). The UV-DDB complex may recognize UV-induced DNA damage and recruit proteins of the nucleotide excision repair pathway (the NER pathway) to initiate DNA repair.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

DDB2 Antibody - Protein Information

Name DDB2

Function

Protein, which is both involved in DNA repair and protein ubiquitination, as part of the UV-DDB complex and DCX (DDB1-CUL4-X-box) complexes, respectively (PubMed:10882109, PubMed:11278856, PubMed:11705987, PubMed:12732143, PubMed:15882621, PubMed:16473935, PubMed:18593899, PubMed:<a

[32789493](http://www.uniprot.org/citations/32789493), PubMed: [9892649](http://www.uniprot.org/citations/9892649)). Core component of the UV-DDB complex (UV-damaged DNA-binding protein complex), a complex that recognizes UV-induced DNA damage and recruit proteins of the nucleotide excision repair pathway (the NER pathway) to initiate DNA repair (PubMed: [10882109](http://www.uniprot.org/citations/10882109), PubMed: [11278856](http://www.uniprot.org/citations/11278856), PubMed: [11705987](http://www.uniprot.org/citations/11705987), PubMed: [12944386](http://www.uniprot.org/citations/12944386), PubMed: [14751237](http://www.uniprot.org/citations/14751237), PubMed: [16260596](http://www.uniprot.org/citations/16260596), PubMed: [32789493](http://www.uniprot.org/citations/32789493)). The UV-DDB complex preferentially binds to cyclobutane pyrimidine dimers (CPD), 6-4 photoproducts (6-4 PP), apurinic sites and short mismatches (PubMed: [10882109](http://www.uniprot.org/citations/10882109), PubMed: [11278856](http://www.uniprot.org/citations/11278856), PubMed: [11705987](http://www.uniprot.org/citations/11705987), PubMed: [12944386](http://www.uniprot.org/citations/12944386), PubMed: [16260596](http://www.uniprot.org/citations/16260596)). Also functions as the substrate recognition module for the DCX (DDB2-CUL4-X-box) E3 ubiquitin-protein ligase complex DDB2-CUL4-ROC1 (also known as CUL4-DDB-ROC1 and CUL4-DDB-RBX1) (PubMed: [12732143](http://www.uniprot.org/citations/12732143), PubMed: [15882621](http://www.uniprot.org/citations/15882621), PubMed: [16473935](http://www.uniprot.org/citations/16473935), PubMed: [18593899](http://www.uniprot.org/citations/18593899), PubMed: [26572825](http://www.uniprot.org/citations/26572825)). The DDB2-CUL4-ROC1 complex may ubiquitinate histone H2A, histone H3 and histone H4 at sites of UV-induced DNA damage (PubMed: [16473935](http://www.uniprot.org/citations/16473935), PubMed: [16678110](http://www.uniprot.org/citations/16678110)). The ubiquitination of histones may facilitate their removal from the nucleosome and promote subsequent DNA repair (PubMed: [16473935](http://www.uniprot.org/citations/16473935), PubMed: [16678110](http://www.uniprot.org/citations/16678110)). The DDB2-CUL4-ROC1 complex also ubiquitinates XPC, which may enhance DNA-binding by XPC and promote NER (PubMed: [15882621](http://www.uniprot.org/citations/15882621)). The DDB2-CUL4-ROC1 complex also ubiquitinates KAT7/HBO1 in response to DNA damage, leading to its degradation: recognizes KAT7/HBO1 following phosphorylation by ATR (PubMed: [26572825](http://www.uniprot.org/citations/26572825)).

Cellular Location

Nucleus. Chromosome. Note=Accumulates at sites of DNA damage following UV irradiation.

Tissue Location

Ubiquitously expressed; with highest levels in corneal endothelium and lowest levels in brain. Isoform D1 is highly expressed in brain and heart. Isoform D2, isoform D3 and isoform D4 are weakly expressed.


DDB2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DDB2 Antibody - Images



Western blot analysis of DDB2 expression in HeLa cell lysate.