

PRMT6 Antibody
Rabbit mAb
Catalog # AP92127

Specification

PRMT6 Antibody - Product Information

Application	WB
Primary Accession	O96LA8
Reactivity	Rat
Clonality	Monoclonal
Other Names	
HRMT1L6; PRMT 6;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	41938 Da

PRMT6 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human PRMT6
Description	Arginine methyltransferase that can catalyze the formation of both omega-N monomethylarginine (MMA) and asymmetrical dimethylarginine (aDMA), with a strong preference for the formation of aDMA. Preferentially methylates arginyl residues present in a glycine and arginine-rich domain and displays preference for monomethylated substrates.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

PRMT6 Antibody - Protein Information

Name PRMT6

Synonyms HRMT1L6

Function

Arginine methyltransferase that can catalyze the formation of both omega-N monomethylarginine (MMA) and asymmetrical dimethylarginine (aDMA), with a strong preference for the formation of aDMA (PubMed:17898714, PubMed:<a href="http://www.uniprot.org/citations/18077460"

target="_blank">18077460, PubMed:18079182, PubMed:19405910, PubMed:30420520). Preferentially methylates arginyl residues present in a glycine and arginine-rich domain and displays preference for monomethylated substrates (PubMed:17898714, PubMed:18077460, PubMed:18079182, PubMed:19405910). Specifically mediates the asymmetric dimethylation of histone H3 'Arg-2' to form H3R2me2a (PubMed:17898714, PubMed:18077460, PubMed:18079182). H3R2me2a represents a specific tag for epigenetic transcriptional repression and is mutually exclusive with methylation on histone H3 'Lys-4' (H3K4me2 and H3K4me3) (PubMed:17898714, PubMed:18077460). Acts as a transcriptional repressor of various genes such as HOXA2, THBS1 and TP53 (PubMed:19509293). Repression of TP53 blocks cellular senescence (By similarity). Also methylates histone H2A and H4 'Arg-3' (H2AR3me and H4R3me, respectively). Acts as a regulator of DNA base excision during DNA repair by mediating the methylation of DNA polymerase beta (POLB), leading to the stimulation of its polymerase activity by enhancing DNA binding and processivity (PubMed:16600869). Methylates HMGA1 (PubMed:16157300, PubMed:16159886). Regulates alternative splicing events. Acts as a transcriptional coactivator of a number of steroid hormone receptors including ESR1, ESR2, PGR and NR3C1. Promotes fasting-induced transcriptional activation of the gluconeogenic program through methylation of the CRTC2 transcription coactivator (By similarity). May play a role in innate immunity against HIV-1 in case of infection by methylating and impairing the function of various HIV-1 proteins such as Tat, Rev and Nucleocapsid protein p7 (NC) (PubMed:17267505). Methylates GPS2, protecting GPS2 from ubiquitination and degradation (By similarity). Methylates SIRT7, inhibiting SIRT7 histone deacetylase activity and promoting mitochondria biogenesis (PubMed:30420520).

Cellular Location

Nucleus.


Tissue Location

Highly expressed in kidney and testis.

PRMT6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PRMT6 Antibody - Images

Western blot analysis of PRMT6 expression in (1) Jurkat cell lysate; (2) NIH/3T3 cell lysate; (3) C6 cell lysate.