

**EHMT2/G9A Antibody**  
Rabbit mAb  
Catalog # AP92247

## Specification

---

### EHMT2/G9A Antibody - Product Information

Application	WB, IHC
Primary Accession	<a href="#">O96KQ7</a>
Clonality	Monoclonal
<b>Other Names</b>	
Bat8; Ehmt2; G9A; GAT8; NG36;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	132370 Da

### EHMT2/G9A Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human EHMT2/G9A
Description	Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

### EHMT2/G9A Antibody - Protein Information

**Name** EHMT2

**Synonyms** BAT8, C6orf30, G9A, KMT1C, NG36

#### Function

Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like

E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself.

#### Cellular Location

Nucleus. Chromosome. Note=Associates with euchromatic regions (PubMed:11316813). Does not associate with heterochromatin (PubMed:11316813).

#### Tissue Location


Expressed in all tissues examined, with high levels in fetal liver, thymus, lymph node, spleen and peripheral blood leukocytes and lower level in bone marrow

### EHMT2/G9A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### EHMT2/G9A Antibody - Images



Western blot analysis of EHMT2/G9A expression in HeLa cell lysate.