

ENT2 Antibody
Rabbit mAb
Catalog # AP92291

Specification

ENT2 Antibody - Product Information

Application	WB, IHC, FC, ICC
Primary Accession	Q14542
Reactivity	Rat
Clonality	Monoclonal
Other Names	
DER12; ei-type; HNP36; Nucleoside transporter; SLC29A2;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	50113 Da

ENT2 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human ENT2
Description	Mediates equilibrative transport of purine, pyrimidine nucleosides and the purine base hypoxanthine. Very less sensitive than SLC29A1 to inhibition by nitrobenzylthioinosine (NBMPR), dipyridamole, dilazep and draflazine.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

ENT2 Antibody - Protein Information

Name SLC29A2 ([HGNC:11004](#))

Synonyms DER12, ENT2, HNP36

Function

Bidirectional uniporter involved in the facilitative transport of nucleosides and nucleobases, and contributes to maintaining their cellular homeostasis (PubMed:[10722669](http://www.uniprot.org/citations/10722669), PubMed:[12527552](http://www.uniprot.org/citations/12527552), PubMed:[12590919](http://www.uniprot.org/citations/12590919), PubMed:[16214850](http://www.uniprot.org/citations/16214850), PubMed:[21795683](http://www.uniprot.org/citations/21795683), PubMed:[9396714](http://www.uniprot.org/citations/9396714)),

href="http://www.uniprot.org/citations/9478986" target="_blank">9478986). Functions as a Na(+)-independent, passive transporter (PubMed:9478986). Involved in the transport of nucleosides such as inosine, adenosine, uridine, thymidine, cytidine and guanosine (PubMed:10722669, PubMed:12527552, PubMed:12590919, PubMed:16214850, PubMed:21795683, PubMed:9396714, PubMed:9478986). Also able to transport purine nucleobases (hypoxanthine, adenine, guanine) and pyrimidine nucleobases (thymine, uracil) (PubMed:16214850, PubMed:21795683). Involved in nucleoside transport at basolateral membrane of kidney cells, allowing liver absorption of nucleoside metabolites (PubMed:12527552). Mediates apical nucleoside uptake into Sertoli cells, thereby regulating the transport of nucleosides in testis across the blood-testis-barrier (PubMed:23639800). Mediates both the influx and efflux of hypoxanthine in skeletal muscle microvascular endothelial cells to control the amount of intracellular hypoxanthine available for xanthine oxidase-mediated ROS production (By similarity).

Cellular Location

Apical cell membrane; Multi-pass membrane protein. Basolateral cell membrane; Multi-pass membrane protein. Note=Localized to the apical membrane of Sertoli cells.

Tissue Location

Highly expressed in skeletal muscle (PubMed:9478986). Expressed in liver, lung, placenta, brain, heart, kidney and ovarian tissues (PubMed:9478986). Expressed in testis at the blood-brain-barrier (PubMed:23639800).


ENT2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ENT2 Antibody - Images





Western blot analysis of ENT2 expression in K562 cell lysate.