

Granulin Antibody
Rabbit mAb
Catalog # AP92340**Specification**

Granulin Antibody - Product Information

| | |
|---|-----------------------------|
| Application | WB, IHC, FC, ICC, IP |
| Primary Accession | P28799 |
| Clonality | Monoclonal |
| Other Names | |
| Acrogranin; CLN11; GEP; GP88; Granulins; GRN; PCDGF; PEPI; PGRN; Proepithelin; Progranulin; | |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 63544 Da |

Granulin Antibody - Additional Information

| | |
|------------------------------|--|
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human Granulin |
| Description | Granulins have possible cytokine-like activity. They may play a role in inflammation, wound repair, and tissue remodeling. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Granulin Antibody - Protein Information**Name** GRN ([HGNC:4601](#))**Function**

Secreted protein that acts as a key regulator of lysosomal function and as a growth factor involved in inflammation, wound healing and cell proliferation (PubMed:[12526812](http://www.uniprot.org/citations/12526812), PubMed:[18378771](http://www.uniprot.org/citations/18378771), PubMed:[28073925](http://www.uniprot.org/citations/28073925), PubMed:[28453791](http://www.uniprot.org/citations/28453791), PubMed:[28541286](http://www.uniprot.org/citations/28541286)). Regulates protein trafficking to lysosomes and, also the activity of lysosomal enzymes (PubMed:[28453791](http://www.uniprot.org/citations/28453791), PubMed:[28541286](http://www.uniprot.org/citations/28541286)). Facilitates also the acidification of lysosomes, causing degradation of mature CTSD by CTSB (PubMed:[28073925](http://www.uniprot.org/citations/28073925)). In addition, functions as a wound-related growth factor that acts directly on dermal fibroblasts and endothelial

cells to promote division, migration and the formation of capillary-like tubule structures (By similarity). Also promotes epithelial cell proliferation by blocking TNF-mediated neutrophil activation preventing release of oxidants and proteases (PubMed:12526812). Moreover, modulates inflammation in neurons by preserving neurons survival, axonal outgrowth and neuronal integrity (PubMed:18378771).

Cellular Location

Secreted. Lysosome Note=Endocytosed by SORT1 and delivered to lysosomes (PubMed:21092856, PubMed:28073925). Targeted to lysosome by PSAP via M6PR and LRP1, in both biosynthetic and endocytic pathways (PubMed:26370502, PubMed:28073925). Co-localized with GBA1 in the intracellular trafficking compartments until to lysosome (By similarity) {ECO:0000250|UniProtKB:P28798, ECO:0000269|PubMed:21092856, ECO:0000269|PubMed:26370502, ECO:0000269|PubMed:28073925}

Tissue Location


In myelogenous leukemic cell lines of promonocytic, promyelocytic, and proerythroid lineage, in fibroblasts, and very strongly in epithelial cell lines. Present in inflammatory cells and bone marrow. Highest levels in kidney

Granulin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Granulin Antibody - Images



Western blot analysis of Granulin expression in 293T cell lysate.