

ADA Antibody
Rabbit mAb
Catalog # AP92392

Specification

ADA Antibody - Product Information

Application	WB, ICC, IP
Primary Accession	P00813
Reactivity	Rat
Clonality	Monoclonal
Other Names	
ada; ADA1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	40764 Da

ADA Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human ADA
Description	Catalyzes the hydrolytic deamination of adenosine and 2-deoxyadenosine. Plays an important role in purine metabolism and in adenosine homeostasis. Modulates signaling by extracellular adenosine, and so contributes indirectly to cellular signaling events.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

ADA Antibody - Protein Information

Name ADA

Synonyms ADA1

Function

Catalyzes the hydrolytic deamination of adenosine and 2- deoxyadenosine (PubMed:16670267, PubMed:23193172, PubMed:26166670, PubMed:8452534, PubMed:9361033). Plays an important role in purine metabolism and in adenosine homeostasis. Modulates signaling by

extracellular adenosine, and so contributes indirectly to cellular signaling events. Acts as a positive regulator of T-cell coactivation, by binding DPP4 (PubMed:20959412). Its interaction with DPP4 regulates lymphocyte-epithelial cell adhesion (PubMed:11772392). Enhances dendritic cell immunogenicity by affecting dendritic cell costimulatory molecule expression and cytokines and chemokines secretion (By similarity). Enhances CD4+ T-cell differentiation and proliferation (PubMed:20959412). Acts as a positive modulator of adenosine receptors ADORA1 and ADORA2A, by enhancing their ligand affinity via conformational change (PubMed:23193172). Stimulates plasminogen activation (PubMed:15016824). Plays a role in male fertility (PubMed:21919946, PubMed:26166670). Plays a protective role in early postimplantation embryonic development (By similarity).

Cellular Location

Cell membrane; Peripheral membrane protein; Extracellular side. Cell junction. Cytoplasmic vesicle lumen {ECO:0000250|UniProtKB:P03958}. Cytoplasm. Lysosome. Note=Colocalized with DPP4 at the cell surface.

Tissue Location

Found in all tissues, occurs in large amounts in T- lymphocytes (PubMed:20959412). Expressed at the time of weaning in gastrointestinal tissues.


ADA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADA Antibody - Images





Western blot analysis of ADA expression in Jurkat cell lysate.