

**MICA Antibody**  
Rabbit mAb  
Catalog # AP92582

## Specification

---

### MICA Antibody - Product Information

|                    |                        |
|--------------------|------------------------|
| Application        | WB                     |
| Primary Accession  | <a href="#">Q29983</a> |
| Clonality          | Monoclonal             |
| <b>Other Names</b> |                        |
| PERB11.1; MICA;    |                        |
| Isotype            | Rabbit IgG             |
| Host               | Rabbit                 |
| Calculated MW      | 42915 Da               |

### MICA Antibody - Additional Information

|                              |  |
|------------------------------|--|
| Purification                 | Affinity-chromatography  |
| Immunogen                    | A synthesized peptide derived from human MICA  |
| Description                  | Seems to have no role in antigen presentation. Acts as a stress-induced self-antigen that is recognized by gamma delta T-cells. Ligand for the KLRK1/NKG2D receptor. Binding to KLRK1 leads to cell lysis. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.                          |

### MICA Antibody - Protein Information

**Name** MICA {ECO:0000312|EMBL:CAI41907.1}

#### Function

Widely expressed membrane-bound protein which acts as a ligand to stimulate an activating receptor KLRK1/NKG2D, expressed on the surface of essentially all human natural killer (NK), gammadelta T and CD8 alphabeta T-cells (PubMed:<a href="http://www.uniprot.org/citations/11777960" target="\_blank">11777960</a>, PubMed:<a href="http://www.uniprot.org/citations/11491531" target="\_blank">11491531</a>). Up-regulated in stressed conditions, such as viral and bacterial infections or DNA damage response, serves as signal of cellular stress, and engagement of KLRK1/NKG2D by MICA triggers NK-cells resulting in a range of immune effector functions, such as cytotoxicity and cytokine production (PubMed:<a href="http://www.uniprot.org/citations/10426993" target="\_blank">10426993</a>).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein. Cytoplasm Note=Expressed on the cell surface in gastric epithelium, endothelial cells and fibroblasts and in the cytoplasm in keratinocytes and monocytes. Infection with human adenovirus 5 suppresses cell surface expression due to the adenoviral E3-19K protein which causes retention in the endoplasmic reticulum.

#### Tissue Location


Widely expressed with the exception of the central nervous system where it is absent. Expressed predominantly in gastric epithelium and also in monocytes, keratinocytes, endothelial cells, fibroblasts and in the outer layer of Hassal's corpuscles within the medulla of normal thymus. In skin, expressed mainly in the keratin layers, basal cells, ducts and follicles. Also expressed in many, but not all, epithelial tumors of lung, breast, kidney, ovary, prostate and colon. In thymomas, overexpressed in cortical and medullar epithelial cells. Tumors expressing MICA display increased levels of gamma delta T-cells.

#### MICA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### MICA Antibody - Images



Western blot analysis of MICA expression in A431 cell lysate.