

**NLK Antibody**  
Rabbit mAb  
Catalog # AP92601

## Specification

---

### NLK Antibody - Product Information

Application	WB, FC
Primary Accession	<a href="#">O9UBE8</a>
Clonality	Monoclonal
<b>Other Names</b>	
LAK1; Nemo like kinase; Nlk;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	58283 Da

### NLK Antibody - Additional Information

Purification	<b>Affinity-chromatography</b>
Immunogen	<b>A synthesized peptide derived from human NLK</b>
Description	<b>Role in cell fate determination, required for differentiation of bone marrow stromal cells. Acts downstream of MAP3K7 and HIPK2 to negatively regulate the canonical Wnt/beta-catenin signaling pathway and the phosphorylation and destruction of the MYB transcription factor.</b>
Storage Condition and Buffer	<b>Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.</b>

### NLK Antibody - Protein Information

**Name** NLK

**Synonyms** LAK1 {ECO:0000312|EMBL:AAD56013.1}

#### Function

Serine/threonine-protein kinase that regulates a number of transcription factors with key roles in cell fate determination (PubMed: [12482967](http://www.uniprot.org/citations/12482967) target="\_blank">12482967</a>, PubMed: [14960582](http://www.uniprot.org/citations/14960582) target="\_blank">14960582</a>, PubMed: [15004007](http://www.uniprot.org/citations/15004007) target="\_blank">15004007</a>, PubMed: [15764709](http://www.uniprot.org/citations/15764709) target="\_blank">15764709</a>, PubMed: [20061393](http://www.uniprot.org/citations/20061393) target="\_blank">20061393</a>, PubMed: [20874444](http://www.uniprot.org/citations/20874444) target="\_blank">20874444</a>, PubMed: [21454679](http://www.uniprot.org/citations/21454679) target="\_blank">21454679</a>)

target="\_blank">21454679</a>). Positive effector of the non-canonical Wnt signaling pathway, acting downstream of WNT5A, MAP3K7/TAK1 and HIPK2 (PubMed:<a href="http://www.uniprot.org/citations/15004007" target="\_blank">15004007</a>, PubMed:<a href="http://www.uniprot.org/citations/15764709" target="\_blank">15764709</a>). Negative regulator of the canonical Wnt/beta-catenin signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/12482967" target="\_blank">12482967</a>). Binds to and phosphorylates TCF7L2/TCF4 and LEF1, promoting the dissociation of the TCF7L2/LEF1/beta-catenin complex from DNA, as well as the ubiquitination and subsequent proteolysis of LEF1 (PubMed:<a href="http://www.uniprot.org/citations/21454679" target="\_blank">21454679</a>). Together these effects inhibit the transcriptional activation of canonical Wnt/beta-catenin target genes (PubMed:<a href="http://www.uniprot.org/citations/12482967" target="\_blank">12482967</a>, PubMed:<a href="http://www.uniprot.org/citations/21454679" target="\_blank">21454679</a>). Negative regulator of the Notch signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/20118921" target="\_blank">20118921</a>). Binds to and phosphorylates NOTCH1, thereby preventing the formation of a transcriptionally active ternary complex of NOTCH1, RBPJ/RBPSUH and MAML1 (PubMed:<a href="http://www.uniprot.org/citations/20118921" target="\_blank">20118921</a>). Negative regulator of the MYB family of transcription factors (PubMed:<a href="http://www.uniprot.org/citations/15082531" target="\_blank">15082531</a>). Phosphorylation of MYB leads to its subsequent proteolysis while phosphorylation of MYBL1 and MYBL2 inhibits their interaction with the coactivator CREBBP (PubMed:<a href="http://www.uniprot.org/citations/15082531" target="\_blank">15082531</a>). Other transcription factors may also be inhibited by direct phosphorylation of CREBBP itself (PubMed:<a href="http://www.uniprot.org/citations/15082531" target="\_blank">15082531</a>). Acts downstream of IL6 and MAP3K7/TAK1 to phosphorylate STAT3, which is in turn required for activation of NLK by MAP3K7/TAK1 (PubMed:<a href="http://www.uniprot.org/citations/15004007" target="\_blank">15004007</a>, PubMed:<a href="http://www.uniprot.org/citations/15764709" target="\_blank">15764709</a>). Upon IL1B stimulus, cooperates with ATF5 to activate the transactivation activity of C/EBP subfamily members (PubMed:<a href="http://www.uniprot.org/citations/25512613" target="\_blank">25512613</a>). Phosphorylates ATF5 but also stabilizes ATF5 protein levels in a kinase-independent manner (PubMed:<a href="http://www.uniprot.org/citations/25512613" target="\_blank">25512613</a>). Acts as an inhibitor of the mTORC1 complex in response to osmotic stress by mediating phosphorylation of RPTOR, thereby preventing recruitment of the mTORC1 complex to lysosomes (PubMed:<a href="http://www.uniprot.org/citations/26588989" target="\_blank">26588989</a>).

#### Cellular Location


Nucleus {ECO:0000250|UniProtKB:O54949}. Cytoplasm {ECO:0000250|UniProtKB:O54949}. Note=Predominantly nuclear. A smaller fraction is cytoplasmic. {ECO:0000250|UniProtKB:O54949}

#### NLK Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### NLK Antibody - Images



Western blot analysis of NLK expression in A375 cell lysate.