

NME2 Antibody
Rabbit mAb
Catalog # AP92613

Specification

NME2 Antibody - Product Information

Application	WB, IHC, FC, ICC
Primary Accession	P22392
Reactivity	Rat
Clonality	Monoclonal

Other Names

C myc purine binding transcription factor PUF; MGC2212; NDKB; NDP kinase B; NDPKB; NM23B; Nucleoside diphosphate kinase B; PUF;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	17298 Da

NME2 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human NME2
Description	Major role in the synthesis of nucleoside triphosphates other than ATP. Negatively regulates Rho activity by interacting with AKAP13/LBC.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

NME2 Antibody - Protein Information

Name NME2

Synonyms NM23B

Function

Major role in the synthesis of nucleoside triphosphates other than ATP. The ATP gamma phosphate is transferred to the NDP beta phosphate via a ping-pong mechanism, using a phosphorylated active-site intermediate (By similarity). Negatively regulates Rho activity by interacting with AKAP13/LBC (PubMed: [15249197](http://www.uniprot.org/citations/15249197)). Acts as a transcriptional activator of the MYC gene; binds DNA non-specifically (PubMed: [19435876](http://www.uniprot.org/citations/19435876), PubMed: [8392752](http://www.uniprot.org/citations/8392752)). Binds to both single-stranded guanine- and cytosine-rich strands within the nuclease hypersensitive element (NHE) III(1) region of the MYC gene promoter. Does not

bind to duplex NHE III(1) (PubMed:19435876). Has G-quadruplex (G4) DNA-binding activity, which is independent of its nucleotide-binding and kinase activity. Binds both folded and unfolded G4 with similar low nanomolar affinities. Stabilizes folded G4s regardless of whether they are prefolded or not (PubMed:25679041). Exhibits histidine protein kinase activity (PubMed:20946858).

Cellular Location

Cytoplasm. Cell projection, lamellipodium. Cell projection, ruffle. Note=Colocalizes with ITGB1 and ITGB1BP1 at the edge or peripheral ruffles and lamellipodia during the early stages of cell spreading on fibronectin or collagen but not on vitronectin or laminin substrates [Isoform 3]: Cytoplasm. Cytoplasm, perinuclear region. Nucleus

Tissue Location


[Isoform 1]: Ubiquitously expressed.

NME2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NME2 Antibody - Images



Western blot analysis of NME2 expression in LnCaP cell lysate.