

## Phospho-Histone H3.3 (S31) Antibody

Rabbit mAb Catalog # AP92817

#### **Specification**

### Phospho-Histone H3.3 (S31) Antibody - Product Information

Application WB, IHC, FC, ICC

Primary Accession P84243
Reactivity Rat

Clonality Monoclonal

Other Names Histone H3.3;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 15328 Da

## Phospho-Histone H3.3 (S31) Antibody - Additional Information

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

**Phospho-Histone H3.3 (S31)** 

Description Variant histone H3 which replaces

conventional H3 in a wide range of nucleosomes in active genes. Constitutes

nucleosomes in active genes. Constitutes the predominant form of histone H3 in non-dividing cells and is incorporated into chromatin independently of DNA synthesis. Rabbit IgG in phosphate buffered saline,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

# Phospho-Histone H3.3 (S31) Antibody - Protein Information

Name H3-3A (<u>HGNC:4764</u>)

Storage Condition and Buffer

Synonyms H3.3A, H3F3, H3F3A

#### **Function**

Variant histone H3 which replaces conventional H3 in a wide range of nucleosomes in active genes. Constitutes the predominant form of histone H3 in non-dividing cells and is incorporated into chromatin independently of DNA synthesis. Deposited at sites of nucleosomal displacement throughout transcribed genes, suggesting that it represents an epigenetic imprint of transcriptionally active chromatin. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of



histones, also called histone code, and nucleosome remodeling.


**Cellular Location** Nucleus. Chromosome

## Phospho-Histone H3.3 (S31) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# Phospho-Histone H3.3 (S31) Antibody - Images



Western blot analysis of Phospho-Histone H3.3 (S31) expression in HeLa cell treated with Calyculin A lysate.