

ZHX2 Antibody

Rabbit mAb Catalog # AP92856

Specification

ZHX2 Antibody - Product Information

Application WB, FC, ICC
Primary Accession Q9Y6X8
Reactivity Rat
Clonality Monoclonal

Other Names

AFP regulator 1; AFR1; Alpha fetoprotein regulator 1; RAF; Regulator of AFP; Transcription factor ZHX2; ZHX2; Zinc finger and homeodomain protein 2; Zinc fingers and homeoboxes 2; Zinc fingers and homeoboxes protein 2;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 92307 Da

ZHX2 Antibody - Additional Information

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

ZHX2

Description Acts as a transcriptional repressor.

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

ZHX2 Antibody - Protein Information

Name ZHX2

Synonyms AFR1, KIAA0854, RAF

Storage Condition and Buffer

Function

Acts as a transcriptional repressor (PubMed:12741956). Represses the promoter activity of the CDC25C gene stimulated by NFYA (PubMed:12741956). May play a role in retinal development where it regulates the composition of bipolar cell populations, by promoting differentiation of bipolar OFF-type cells (By similarity). In the brain, may promote maintenance and suppress differentiation of neural progenitor cells in the developing cortex (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:12741956, ECO:0000269|PubMed:17056598} Note=Colocalizes with EFNB1 intracellular domain in the



nucleus {ECO:0000250|UniProtKB:Q8C0C0}

Tissue Location


Ubiquitously expressed. Expressed in podocytes.

ZHX2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ZHX2 Antibody - Images



Western blot analysis of ZHX2 expression in HeLa cell lysate.