

CD223 Antibody
Rabbit mAb
Catalog # AP93162**Specification**

CD223 Antibody - Product Information

Application	IHC
Primary Accession	P18627
Clonality	Monoclonal
Other Names	
CD223; Lag3; Lymphocyte activating 3; Protein FDC;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	57449 Da

CD223 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CD223
Description	Involved in lymphocyte activation. Binds to HLA class-II antigens.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

CD223 Antibody - Protein Information**Name** LAG3 ([HGNC:6476](#))**Synonyms** FDC**Function**

Lymphocyte activation gene 3 protein: Inhibitory receptor on antigen activated T-cells (PubMed:20421648, PubMed:7805750, PubMed:8647185). Delivers inhibitory signals upon binding to ligands, such as FGL1 (By similarity). FGL1 constitutes a major ligand of LAG3 and is responsible for LAG3 T-cell inhibitory function (By similarity). Following TCR engagement, LAG3 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1, possibly by acting as a coreceptor for PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed:20421648, PubMed:7805750, PubMed:8647185).

target="_blank">8647185). Also mediates immune tolerance: constitutively expressed on a subset of regulatory T-cells (Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity). Binds MHC class II (MHC-II); the precise role of MHC-II-binding is however unclear (PubMed:8647185).

Cellular Location

[Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein

Tissue Location

Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.

CD223 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD223 Antibody - Images