

**Phospho-GLUT4 (S488) Antibody**  
Rabbit mAb  
Catalog # AP93244

**Specification**

---

**Phospho-GLUT4 (S488) Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P14672</a>
Reactivity	Rat
Clonality	Monoclonal

**Other Names**

insulin-responsive; Glucose transporter GLUT 4; Glucose transporter type 4; Glucose transporter type 4 insulin responsive; GLUT4; Insulin responsive glucose transporter type 4; SLC2A4 ; Solute carrier family 2 member 4;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	54787 Da

**Phospho-GLUT4 (S488) Antibody - Additional Information**

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Phospho-GLUT4 (S488)
Description	Insulin-regulated facilitative glucose transporter.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**Phospho-GLUT4 (S488) Antibody - Protein Information**

Name SLC2A4 ([HGNC:11009](#))

**Function**

Insulin-regulated facilitative glucose transporter, which plays a key role in removal of glucose from circulation. Response to insulin is regulated by its intracellular localization: in the absence of insulin, it is efficiently retained intracellularly within storage compartments in muscle and fat cells. Upon insulin stimulation, translocates from these compartments to the cell surface where it transports glucose from the extracellular milieu into the cell.

**Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:P14142}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P14142} Endomembrane system; Multi-pass membrane protein. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:P14142}. Note=Localizes primarily to the perinuclear region, undergoing continued recycling to the plasma membrane where it is rapidly reinternalized (PubMed:8300557). The dileucine internalization motif is critical for intracellular

sequestration (PubMed:8300557). Insulin stimulation induces translocation to the cell membrane (By similarity) {ECO:0000250|UniProtKB:P14142, ECO:0000269|PubMed:8300557}

**Tissue Location**

Skeletal and cardiac muscles; brown and white fat.

**Phospho-GLUT4 (S488) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Phospho-GLUT4 (S488) Antibody - Images**