

SYVN1 / HRD1 Antibody
Rabbit mAb
Catalog # AP93248**Specification****SYVN1 / HRD1 Antibody - Product Information**

Application	WB, FC, ICC, IP
Primary Accession	Q86TM6
Reactivity	Rat
Clonality	Monoclonal

Other Names

DER3; E3 ubiquitin-protein ligase synoviolin; HRD1; Synovial apoptosis inhibitor 1; Synoviolin 1; Synoviolin 1 isoform b; SYNOVIOLIN; SYVN1;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	67685 Da

SYVN1 / HRD1 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SYVN1 / HRD1
Description	Acts as an E3 ubiquitin-protein ligase which accepts ubiquitin specifically from endoplasmic reticulum-associated UBC7 E2 ligase and transfers it to substrates, promoting their degradation.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

SYVN1 / HRD1 Antibody - Protein Information

Name SYVN1

Synonyms HRD1, KIAA1810

Function

E3 ubiquitin-protein ligase which accepts ubiquitin specifically from endoplasmic reticulum-associated UBC7 E2 ligase and transfers it to substrates, promoting their degradation (PubMed: [12459480](http://www.uniprot.org/citations/12459480), PubMed: [12646171](http://www.uniprot.org/citations/12646171), PubMed: [12975321](http://www.uniprot.org/citations/12975321), PubMed: [14593114](http://www.uniprot.org/citations/14593114), PubMed: [16289116](http://www.uniprot.org/citations/16289116), PubMed: [16847254](http://www.uniprot.org/citations/16847254)),

PubMed: 17059562, PubMed: 17141218, PubMed: 17170702, PubMed: 22607976, PubMed: 26471130, PubMed: 28827405). Component of the endoplasmic reticulum quality control (ERQC) system also called ER-associated degradation (ERAD) involved in ubiquitin- dependent degradation of misfolded endoplasmic reticulum proteins (PubMed: 12459480, PubMed: 12646171, PubMed: 12975321, PubMed: 14593114, PubMed: 16289116, PubMed: 16847254, PubMed: 17059562, PubMed: 17141218, PubMed: 17170702, PubMed: 22607976, PubMed: 26471130, PubMed: 28842558). Also promotes the degradation of normal but naturally short-lived proteins such as SGK. Protects cells from ER stress-induced apoptosis. Protects neurons from apoptosis induced by polyglutamine-expanded huntingtin (HTT) or unfolded GPR37 by promoting their degradation (PubMed: 17141218). Sequesters p53/TP53 in the cytoplasm and promotes its degradation, thereby negatively regulating its biological function in transcription, cell cycle regulation and apoptosis (PubMed: 17170702). Mediates the ubiquitination and subsequent degradation of cytoplasmic NFE2L1 (By similarity). During the early stage of B cell development, required for degradation of the pre-B cell receptor (pre-BCR) complex, hence supporting further differentiation into mature B cells (By similarity).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein

Tissue Location

Ubiquitously expressed, with highest levels in liver and kidney (at protein level). Up-regulated in synovial tissues from patients with rheumatoid arthritis (at protein level)

SYVN1 / HRD1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SYVN1 / HRD1 Antibody - Images