

**Phospho-PRAS40 (T246) Antibody**  
Rabbit mAb  
Catalog # AP93278**Specification****Phospho-PRAS40 (T246) Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O96B36</a>
Clonality	Monoclonal
<b>Other Names</b>	
AKT1S1; PRAS; PRAS40;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	27383 Da

**Phospho-PRAS40 (T246) Antibody - Additional Information**

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Phospho-PRAS40 (T246)
Description	Subunit of mTORC1, which regulates cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**Phospho-PRAS40 (T246) Antibody - Protein Information**

**Name** AKT1S1 {ECO:0000312|EMBL:AAH16043.1}

**Function**

Negative regulator of the mechanistic target of rapamycin complex 1 (mTORC1), an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:<a href="http://www.uniprot.org/citations/17277771" target="\_blank">17277771</a>, PubMed:<a href="http://www.uniprot.org/citations/17386266" target="\_blank">17386266</a>, PubMed:<a href="http://www.uniprot.org/citations/17510057" target="\_blank">17510057</a>, PubMed:<a href="http://www.uniprot.org/citations/29236692" target="\_blank">29236692</a>). In absence of insulin and nutrients, AKT1S1 associates with the mTORC1 complex and directly inhibits mTORC1 activity by blocking the MTOR substrate- recruitment site (PubMed:<a href="http://www.uniprot.org/citations/29236692" target="\_blank">29236692</a>). In response to insulin and nutrients, AKT1S1 dissociates from mTORC1 (PubMed:<a href="http://www.uniprot.org/citations/17386266" target="\_blank">17386266</a>, PubMed:<a href="http://www.uniprot.org/citations/17386266" target="\_blank">17386266</a>, PubMed:<a href="http://www.uniprot.org/citations/17386266" target="\_blank">17386266</a>).

href="http://www.uniprot.org/citations/18372248" target="\_blank">18372248</a>). Its activity is dependent on its phosphorylation state and binding to 14-3-3 (PubMed:<a href="http://www.uniprot.org/citations/16174443" target="\_blank">16174443</a>, PubMed:<a href="http://www.uniprot.org/citations/18372248" target="\_blank">18372248</a>). May also play a role in nerve growth factor-mediated neuroprotection (By similarity).

**Cellular Location**

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9D1F4}. Note=Found in the cytosolic fraction of the brain. {ECO:0000250|UniProtKB:Q9D1F4}

**Tissue Location**

Widely expressed with highest levels of expression in liver and heart. Expressed at higher levels in cancer cell lines (e.g. A-549 and HeLa) than in normal cell lines (e.g. HEK293)

**Phospho-PRAS40 (T246) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Phospho-PRAS40 (T246) Antibody - Images**