

MOTS-C Rabbit Polyclonal Antibody
MOTS-C Rabbit Polyclonal Antibody
Catalog # AP93622**Specification**

MOTS-C Rabbit Polyclonal Antibody - Product Information

Application	IHC
Primary Accession	AOA0C5B5G6
Reactivity	Rat, Human, Mouse
Clonality	Polyclonal
Calculated MW	2175

MOTS-C Rabbit Polyclonal Antibody - Additional Information**Other Names**

Mitochondrial-derived peptide MOTS-c, Mitochondrial open reading frame of the 12S rRNA-c, MT-RNR1 ([HGNC:7470](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=7470))

Storage Conditions

-20°C

MOTS-C Rabbit Polyclonal Antibody - Protein Information

Name MT-RNR1 ([HGNC:7470](#))

Function

Regulates insulin sensitivity and metabolic homeostasis (PubMed:[25738459](http://www.uniprot.org/citations/25738459), PubMed:[33468709](http://www.uniprot.org/citations/33468709)). Inhibits the folate cycle, thereby reducing de novo purine biosynthesis which leads to the accumulation of the de novo purine synthesis intermediate 5-aminoimidazole-4- carboxamide (AICAR) and the activation of the metabolic regulator 5'- AMP-activated protein kinase (AMPK) (PubMed:[25738459](http://www.uniprot.org/citations/25738459)). Protects against age-dependent and diet-induced insulin resistance as well as diet- induced obesity (PubMed:[25738459](http://www.uniprot.org/citations/25738459)). In response to metabolic stress, translocates to the nucleus where it binds to antioxidant response elements (ARE) present in the promoter regions of a number of genes and plays a role in regulating nuclear gene expression in an NFE2L2- dependent manner and increasing cellular resistance to metabolic stress (PubMed:[29983246](http://www.uniprot.org/citations/29983246)). Increases mitochondrial respiration and levels of CPT1A and cytokines IL1B, IL6, IL8, IL10 and TNF in senescent cells (PubMed:[29886458](http://www.uniprot.org/citations/29886458)). Increases activity of the serine/threonine protein kinase complex mTORC2 and reduces activity of the PTEN phosphatase, thus promoting phosphorylation of AKT (PubMed:[33554779](http://www.uniprot.org/citations/33554779)). This promotes AKT-mediated phosphorylation of transcription factor FOXO1 which reduces FOXO1 activity, leading to reduced levels of MSTN and promotion of skeletal muscle growth (PubMed:[33554779](#)).

[33554779](http://www.uniprot.org/citations/33554779)). Promotes osteogenic differentiation of bone marrow mesenchymal stem cells via the TGFB/SMAD pathway (PubMed:[30468456](http://www.uniprot.org/citations/30468456)). Promotes osteoblast proliferation and osteoblast synthesis of type I collagens COL1A1 and COL1A2 via the TGFB/SMAD pathway (PubMed:[31081069](http://www.uniprot.org/citations/31081069)).

Cellular Location

Secreted. Mitochondrion. Nucleus Note=Translocates to the nucleus in response to metabolic stress in an AMPK-dependent manner.

Tissue Location


Detected in plasma (at protein level) (PubMed:25738459, PubMed:32182209). Also expressed in skeletal muscle (at protein level) (PubMed:32182209).

MOTS-C Rabbit Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MOTS-C Rabbit Polyclonal Antibody - Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).