

Chk1 (7N8) Rabbit Monoclonal Antibody
Chk1 (7N8) Rabbit Monoclonal Antibody
Catalog # AP93671**Specification**

Chk1 (7N8) Rabbit Monoclonal Antibody - Product Information

Application	WB, IF, FC, ICC, IP
Primary Accession	O14757
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Calculated MW	54434

Chk1 (7N8) Rabbit Monoclonal Antibody - Additional Information**Gene ID** 1111**Other Names**

Serine/threonine-protein kinase Chk1, 2.7.11.1, CHK1 checkpoint homolog, Cell cycle checkpoint kinase, Checkpoint kinase-1, CHEK1, CHK1

Storage Conditions

-20°C

Chk1 (7N8) Rabbit Monoclonal Antibody - Protein Information**Name** CHEK1**Synonyms** CHK1**Function**

Serine/threonine-protein kinase which is required for checkpoint-mediated cell cycle arrest and activation of DNA repair in response to the presence of DNA damage or unreplicated DNA (PubMed: [11535615](http://www.uniprot.org/citations/11535615), PubMed: [12399544](http://www.uniprot.org/citations/12399544), PubMed: [12446774](http://www.uniprot.org/citations/12446774), PubMed: [14559997](http://www.uniprot.org/citations/14559997), PubMed: [14988723](http://www.uniprot.org/citations/14988723), PubMed: [15311285](http://www.uniprot.org/citations/15311285), PubMed: [15650047](http://www.uniprot.org/citations/15650047), PubMed: [15665856](http://www.uniprot.org/citations/15665856), PubMed: [32357935](http://www.uniprot.org/citations/32357935)). May also negatively regulate cell cycle progression during unperturbed cell cycles (PubMed: [11535615](http://www.uniprot.org/citations/11535615), PubMed: [12399544](http://www.uniprot.org/citations/12399544), PubMed: [12446774](http://www.uniprot.org/citations/12446774), PubMed: [14559997](http://www.uniprot.org/citations/14559997), PubMed: [14988723](http://www.uniprot.org/citations/14988723), PubMed: [15311285](http://www.uniprot.org/citations/15311285), PubMed: [15650047](http://www.uniprot.org/citations/15650047), PubMed: [15665856](http://www.uniprot.org/citations/15665856), PubMed: [32357935](http://www.uniprot.org/citations/32357935)).

href="http://www.uniprot.org/citations/15650047" target="_blank">15650047, PubMed:15665856). This regulation is achieved by a number of mechanisms that together help to preserve the integrity of the genome (PubMed:11535615, PubMed:12399544, PubMed:12446774, PubMed:14559997, PubMed:14988723, PubMed:15311285, PubMed:15650047, PubMed:15665856). Recognizes the substrate consensus sequence [R-X-X- S/T] (PubMed:11535615, PubMed:12399544, PubMed:12446774, PubMed:14559997, PubMed:14988723, PubMed:15311285, PubMed:15650047, PubMed:15665856). Binds to and phosphorylates CDC25A, CDC25B and CDC25C (PubMed:12676583, PubMed:12676925, PubMed:12759351, PubMed:14559997, PubMed:14681206, PubMed:19734889, PubMed:9278511). Phosphorylation of CDC25A at 'Ser-178' and 'Thr-507' and phosphorylation of CDC25C at 'Ser-216' creates binding sites for 14-3-3 proteins which inhibit CDC25A and CDC25C (PubMed:9278511). Phosphorylation of CDC25A at 'Ser-76', 'Ser-124', 'Ser-178', 'Ser-279' and 'Ser-293' promotes proteolysis of CDC25A (PubMed:12676583, PubMed:12676925, PubMed:12759351, PubMed:14681206, PubMed:19734889, PubMed:9278511). Phosphorylation of CDC25A at 'Ser-76' primes the protein for subsequent phosphorylation at 'Ser-79', 'Ser-82' and 'Ser-88' by NEK11, which is required for polyubiquitination and degradation of CDC25A (PubMed:19734889, PubMed:20090422, PubMed:9278511). Inhibition of CDC25 leads to increased inhibitory tyrosine phosphorylation of CDK-cyclin complexes and blocks cell cycle progression (PubMed:9278511). Also phosphorylates NEK6 (PubMed:18728393). Binds to and phosphorylates RAD51 at 'Thr-309', which promotes the release of RAD51 from BRCA2 and enhances the association of RAD51 with chromatin, thereby promoting DNA repair by homologous recombination (PubMed:15665856). Phosphorylates multiple sites within the C-terminus of TP53, which promotes activation of TP53 by acetylation and promotes cell cycle arrest and suppression of cellular proliferation (PubMed:10673501, PubMed:15659650, PubMed:16511572). Also promotes repair of DNA cross-links through

phosphorylation of FANCE (PubMed:17296736). Binds to and phosphorylates TLK1 at 'Ser-743', which prevents the TLK1-dependent phosphorylation of the chromatin assembly factor ASF1A (PubMed:12660173, PubMed:12955071). This may enhance chromatin assembly both in the presence or absence of DNA damage (PubMed:12660173, PubMed:12955071). May also play a role in replication fork maintenance through regulation of PCNA (PubMed:18451105). May regulate the transcription of genes that regulate cell-cycle progression through the phosphorylation of histones (By similarity). Phosphorylates histone H3.1 (to form H3T11ph), which leads to epigenetic inhibition of a subset of genes (By similarity). May also phosphorylate RB1 to promote its interaction with the E2F family of transcription factors and subsequent cell cycle arrest (PubMed:17380128). Phosphorylates SPRTN, promoting SPRTN recruitment to chromatin (PubMed:31316063). Reduces replication stress and activates the G2/M checkpoint, by phosphorylating and inactivating PABIR1/FAM122A and promoting the serine/threonine-protein phosphatase 2A-mediated dephosphorylation and stabilization of WEE1 levels and activity (PubMed:33108758).

Cellular Location

Nucleus. Chromosome. Cytoplasm Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Nuclear export is mediated at least in part by XPO1/CRM1 (PubMed:12676962). Also localizes to the centrosome specifically during interphase, where it may protect centrosomal CDC2 kinase from inappropriate activation by cytoplasmic CDC25B (PubMed:15311285). Proteolytic cleavage at the C-terminus by SPRTN promotes removal from chromatin (PubMed:31316063)

Tissue Location


Expressed ubiquitously with the most abundant expression in thymus, testis, small intestine and colon


Chk1 (7N8) Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Chk1 (7N8) Rabbit Monoclonal Antibody - Images





Western blot detection of Chk1 in K562,3T3,HeLa cell lysates using Chk1 antibody(1:1000 diluted).