

TTLL12 (13V7) Mouse Monoclonal antibody
TTLL12 (13V7) Mouse Monoclonal antibody
Catalog # AP93880**Specification**

TTLL12 (13V7) Mouse Monoclonal antibody - Product Information

Application	WB
Primary Accession	Q14166
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Calculated MW	74404

TTLL12 (13V7) Mouse Monoclonal antibody - Additional Information**Gene ID** 23170**Other Names**

Tubulin--tyrosine ligase-like protein 12, Inactive tubulin--tyrosine ligase-like protein 12, TTLL12, KIAA0153

Storage Conditions

-20°C

TTLL12 (13V7) Mouse Monoclonal antibody - Protein Information**Name** TTLL12**Synonyms** KIAA0153**Function**

Negatively regulates post-translational modifications of tubulin, including detyrosination of the C-terminus and polyglutamylation of glutamate residues (PubMed:[20162578](http://www.uniprot.org/citations/20162578), PubMed:[23251473](http://www.uniprot.org/citations/23251473)). Also, indirectly promotes histone H4 trimethylation at 'Lys-20' (H4K20me3) (PubMed:[23251473](http://www.uniprot.org/citations/23251473)). Probably by controlling tubulin and/or histone H4 post-translational modifications, plays a role in mitosis and in maintaining chromosome number stability (PubMed:[20162578](http://www.uniprot.org/citations/20162578), PubMed:[23251473](http://www.uniprot.org/citations/23251473)). During RNA virus-mediated infection, acts as a negative regulator of the RIG-I pathway by preventing MAVS binding to TBK1 and IKKε (PubMed:[28011935](http://www.uniprot.org/citations/28011935)).

Cellular Location

Cytoplasm. Midbody Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Nucleus Note=Predominantly localizes in the cytoplasm (PubMed:28011935) Partially colocalizes with vimentin in prostate cancer cells

(PubMed:20162578).

Tissue Location


Expressed in the basal layer of prostate and endothelial cells. Increased expression in prostatic intraepithelial neoplasia and metastatic lesions.

TTLL12 (13V7) Mouse Monoclonal antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TTLL12 (13V7) Mouse Monoclonal antibody - Images



HEK293T cells transfected with either overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-TTLL12 antibody (AP93880), and then analyzed by flow cytometry.



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TTLL12 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TTLL12. Positive lysates (100ug) and

(20ug) can be purchased separately from biodragon.