

ASCC2 (5S8) Mouse Monoclonal antibody
ASCC2 (5S8) Mouse Monoclonal antibody
Catalog # AP93895**Specification**

ASCC2 (5S8) Mouse Monoclonal antibody - Product Information

Application	WB, IF
Primary Accession	O9H1I8
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Calculated MW	86360

ASCC2 (5S8) Mouse Monoclonal antibody - Additional Information

Gene ID 84164

Other Names

Activating signal cointegrator 1 complex subunit 2, ASC-1 complex subunit p100, Trip4 complex subunit p100, ASCC2, ASC1P100, RQT3 {ECO:0000303|PubMed:32099016}

Storage Conditions

-20°C

ASCC2 (5S8) Mouse Monoclonal antibody - Protein Information

Name ASCC2

Synonyms ASC1P100, RQT3 {ECO:0000303|PubMed:32099}

Function

Ubiquitin-binding protein involved in DNA repair and rescue of stalled ribosomes (PubMed: 29144457, PubMed: 32099016, PubMed: 32579943, PubMed: 36302773). Plays a role in DNA damage repair as component of the ASCC complex (PubMed: 29144457). Recruits ASCC3 and ALKBH3 to sites of DNA damage by binding to polyubiquitinated proteins that have 'Lys-63'-linked polyubiquitin chains (PubMed: 29144457). Part of the ASC-1 complex that enhances NF-kappa-B, SRF and AP1 transactivation (PubMed: 12077347). Involved in activation of the ribosome quality control (RQC) pathway, a pathway that degrades nascent peptide chains during problematic translation (PubMed: 32099016, PubMed: 32579943, PubMed: 36302773). Specifically recognizes and binds RPS20/uS10 ubiquitinated by ZNF598, promoting recruitment of the RQT (ribosome quality control trigger) complex on stalled ribosomes, followed by disassembly

of stalled ribosomes (PubMed:36302773).

Cellular Location

Nucleus. Nucleus speckle Note=Colocalizes with the spliceosomal components PRPF8 and SNRNP200/BRR2 in nuclear foci when cells have been exposed to alkylating agents that cause DNA damage. Colocalizes with RNF113A and 'Lys-63'-linked polyubiquitinated proteins, ALKBH3 and ASCC3 in nuclear foci when cells have been exposed to alkylating agents that cause DNA damage.

Tissue Location


Ubiquitous..

ASCC2 (5S8) Mouse Monoclonal antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ASCC2 (5S8) Mouse Monoclonal antibody - Images



Anti-ASCC2 mouse monoclonal antibody (AP93895) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ASCC2.



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ASCC2 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ASCC2. Positive lysates (100ug) and (20ug) can be purchased separately from biodragon.