

HAVCR1 Rabbit pAb

HAVCR1 Rabbit pAb Catalog # AP93922

## Specification

# HAVCR1 Rabbit pAb - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW

O5ONS5 Mouse Rabbit Polyclonal 33361

WB

## HAVCR1 Rabbit pAb - Additional Information

Gene ID 171283

**Other Names** 

Hepatitis A virus cellular receptor 1 homolog, HAVcr-1, Kidney injury molecule 1, KIM-1, T cell immunoglobulin and mucin domain-containing protein 1, TIMD-1, T cell membrane protein 1, T-cell immunoglobulin mucin receptor 1, TIM-1, CD365, Havcr1, Kim1, Tim1, Timd1

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## HAVCR1 Rabbit pAb - Protein Information

Name Havcr1

Synonyms Kim1, Tim1, Timd1

#### Function

Phosphatidylserine receptor that plays an important functional role in regulatory B-cells homeostasis including generation, expansion and suppressor functions (PubMed:<a href="http://www.uniprot.org/citations/21821911" target="\_blank">21821911</a>, PubMed:<a href="http://www.uniprot.org/citations/25645598" target="\_blank">25645598</a>, PubMed:<a href="http://www.uniprot.org/citations/25645598" target="\_blank">25645598</a>, PubMed:<a href="http://www.uniprot.org/citations/32668241" target="\_blank">32668241</a>). As P-selectin/SELPLG ligand, plays a specialized role in activated but not naive T-cell trafficking during inflammatory responses (PubMed:<a href="http://www.uniprot.org/citations/24703780" target="\_blank">24703780" target="\_blank">24703780</a>). Controls thereby T-cell accumulation in the inflamed central nervous system (CNS) and the induction of autoimmune disease (PubMed:<a href="http://www.uniprot.org/citations/24703780" target="\_blank">24703780</a>). Regulates also expression of various anti-inflammatory cytokines and co-inhibitory ligands including IL10 (PubMed:<a href="http://www.uniprot.org/citations/25582854" target="\_blank">25582854</a>,



PubMed:<a href="http://www.uniprot.org/citations/25645598" target="\_blank">25645598</a>). Acts as a regulator of T-cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/15793576" target="\_blank">15793576</a>). May play a role in kidney injury and repair (By similarity).

Cellular Location Cell membrane {ECO:0000250|UniProtKB:Q96D42}; Single-pass type I membrane protein


**Tissue Location** Expressed by stimulated T-cells. Expressed during primary antigen stimulation (PubMed:11725301). Expressed at higher levels on B rather than T-cells, both constitutively and after activation (PubMed:21821911).

### HAVCR1 Rabbit pAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## HAVCR1 Rabbit pAb - Images



Sample: Lane 1: Mouse Cerebrum tissue lysates Lane 2: Rat Cerebrum tissue lysates Primary: Anti-HAVCR1 (AP93922) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 39 kDa Observed band size: 50 kDa

## HAVCR1 Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.