

EMT Research Antibody Assembly kit
EMT Research Antibody Assembly kit
Catalog # AP94100**Specification**

EMT Research Antibody Assembly kit - Product Information

Application	IHC-P
Host	Rabbit
Clonality	Polyclonal

EMT Research Antibody Assembly kit - Additional Information**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol


Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

EMT Research Antibody Assembly kit - Protein Information**EMT Research Antibody Assembly kit - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


EMT Research Antibody Assembly kit - Images




Primary: Anti-Occludin (bs-10011R) Paraformaldehyde-fixed, paraffin embedded (Normal skin); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Occludin) Polyclonal Antibody, Unconjugated (bs-10011R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Primary: Anti-N-cadherin (bs-1172R) Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-N-cadherin Polyclonal Antibody, Unconjugated(bs-1172R) 1:400, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Primary: Anti- ZO-1 (bs-1329R) Independently Validated Antibody, image provided by Science Direct, badge number 029577:Formalin-fixed and paraffin embedded human testis labeled with Anti-ZO-1 Polyclonal Antibody, Unconjugated (bs-1329R) at 1:250 followed by conjugation to the secondary antibody.



Primary: Anti- Vimentin (bs-0756R) Paraformaldehyde-fixed, paraffin embedded (human cervical cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Vimentin) Polyclonal Antibody, Unconjugated (bs-0756R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

EMT Research Antibody Assembly kit - Background

For research use only. Not intended for diagnostic or therapeutic use.