

phospho-PI3 Kinase p110 beta (Ser1070) Rabbit pAb
phospho-PI3 Kinase p110 beta (Ser1070) Rabbit pAb
Catalog # AP94133

Specification

phospho-PI3 Kinase p110 beta (Ser1070) Rabbit pAb - Product Information

| | |
|-------------------|------------------------|
| Application | IHC-P, WB |
| Primary Accession | Q24JU2 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |

phospho-PI3 Kinase p110 beta (Ser1070) Rabbit pAb - Additional Information

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


phospho-PI3 Kinase p110 beta (Ser1070) Rabbit pAb - Protein Information


phospho-PI3 Kinase p110 beta (Ser1070) Rabbit pAb - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

phospho-PI3 Kinase p110 beta (Ser1070) Rabbit pAb - Images






Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-phospho-PI3 Kinase p110 beta(Ser1070) Polyclonal Antibody, Unconjugated (AP94133) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Tissue/cell: human colon carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-phospho-PI3 Kinase p110 beta(Ser1070) Polyclonal Antibody, Unconjugated (AP94133) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Tissue/cell: human gastric carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at


37°C for 20 min; Incubation: Anti-phospho-PI3 Kinase p110 beta(Ser1070) Polyclonal Antibody, Unconjugated (AP94133) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-PI3 Kinase p110 beta(Ser1070)) Polyclonal Antibody, Unconjugated (AP94133) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Sample: Lane 1: Human MCF-7 cell lysates Lane 2: Human K562 cell lysates Primary: Anti-phospho-PI3 Kinase p110 beta (Ser1070) (AP94133) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 110-123 kD Observed band size: 110 kD



Paraformaldehyde-fixed, paraffin embedded (rat colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-PI3 Kinase p110 beta (Ser1070)) Polyclonal Antibody, Unconjugated (AP94133) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-PI3 Kinase p110 beta (Ser1070)) Polyclonal Antibody, Unconjugated (AP94133) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

phospho-PI3 Kinase p110 beta (Ser1070) Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.