

HNF4A Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9435a**Specification**

HNF4A Antibody (N-term) - Product Information

Application	IF, WB,E
Primary Accession	P41235
Other Accession	Q9WUU6 , Q14541 , Q91766 , P22449 , P49698
Reactivity	Human
Predicted	Mouse, Rat, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52785
Antigen Region	107-139

HNF4A Antibody (N-term) - Additional Information**Gene ID** 3172**Other Names**

Hepatocyte nuclear factor 4-alpha, HNF-4-alpha, Nuclear receptor subfamily 2 group A member 1, Transcription factor 14, TCF-14, Transcription factor HNF-4, HNF4A, HNF4, NR2A1, TCF14

Target/Specificity

This HNF4A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 107-139 amino acids from the N-terminal region of human HNF4A.

Dilution

IF~~1:10~50

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HNF4A Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HNF4A Antibody (N-term) - Protein Information**Name** HNF4A

Synonyms HNF4, NR2A1, TCF14

Function Transcriptional regulator which controls the expression of hepatic genes during the transition of endodermal cells to hepatic progenitor cells, facilitating the recruitment of RNA pol II to the promoters of target genes (PubMed:[30597922](#)). Activates the transcription of CYP2C38 (By similarity). Represses the CLOCK-BMAL1 transcriptional activity and is essential for circadian rhythm maintenance and period regulation in the liver and colon cells (PubMed:[30530698](#)).

Cellular Location


Nucleus.

HNF4A Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HNF4A Antibody (N-term) - Images



Confocal immunofluorescent analysis of HNF4A Antibody (N-term)(Cat#AP9435a) with MCF-7 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



Anti-HNF4A Antibody (N-term) at 1:1000 dilution + human liver lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 53 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

HNF4A Antibody (N-term) - Background

HNF4A is a nuclear transcription factor which binds DNA as a homodimer. The encoded protein controls the expression of several genes, including hepatocyte nuclear factor 1 alpha, a transcription factor which regulates the expression of several hepatic genes. This gene may play a role in development of the liver, kidney, and intestines.

HNF4A Antibody (N-term) - References

Koh, W., et al. BMC Genomics 11 SUPPL 1, S6 (2010)
Wang, F., et al. Chin. Med. J. 122(20):2477-2482(2009)
Taghavi, S.M., et al. Cardiovasc Diabetol 8, 63 (2009)
Guo, H., et al. Biochem. J. 394 (PT 2), 379-387 (2006)
Hong, Y.H., et al. J. Biol. Chem. 278(30):27495-27501(2003)