

PPAR delta + beta Rabbit pAb
PPAR delta + beta Rabbit pAb
Catalog # AP94559**Specification**

PPAR delta + beta Rabbit pAb - Product Information

Application	WB, IHC-P
Primary Accession	P35396
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49715

PPAR delta + beta Rabbit pAb - Additional Information**Gene ID** 19015**Other Names**

Peroxisome proliferator-activated receptor delta, PPAR-delta, Nuclear hormone receptor 1, NUC1, Nuclear receptor subfamily 1 group C member 2, Peroxisome proliferator-activated receptor beta, PPAR-beta, Ppard, Nr1c2, Pparb

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PPAR delta + beta Rabbit pAb - Protein Information**Name** Ppard**Synonyms** Nr1c2, Pparb**Function**

Ligand-activated transcription factor key mediator of energy metabolism in adipose tissues (PubMed:35675826). Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Has a preference for poly-unsaturated fatty acids, such as gamma- linoleic acid and eicosapentanoic acid. Once activated by a ligand, the receptor binds to promoter elements of target genes. Regulates the peroxisomal beta-oxidation pathway of fatty acids. Functions as transcription activator for the acyl-CoA oxidase gene. Decreases expression of NPC1L1 once activated by a ligand (By similarity).

Cellular Location

Nucleus.


Tissue Location

Heart, adrenal and intestine.


PPAR delta + beta Rabbit pAb - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PPAR delta + beta Rabbit pAb - Images

Sample: Lane 1: Mouse Placenta tissue lysates Lane 2: Mouse Testis tissue lysates Primary: Anti-PPAR delta + beta (AP94559) at 1/200 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 48 kDa Observed band size: 53 kDa



Blank control: A431. Primary Antibody (green line): Rabbit Anti-PPAR delta + beta antibody (AP94559) Dilution: 1 µg /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1 µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Tissue/cell: Human esophageal carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-PPAR-delta Polyclonal Antibody, Unconjugated(AP94559) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

PPAR delta + beta Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.