

**Neuropilin 1 Rabbit pAb**  
**Neuropilin 1 Rabbit pAb**  
**Catalog # AP94626**

**Specification**

---

**Neuropilin 1 Rabbit pAb - Product Information**

Application	IHC-P, WB
Primary Accession	<a href="#">Q5T7F3</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

**Neuropilin 1 Rabbit pAb - Additional Information**

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


**Neuropilin 1 Rabbit pAb - Protein Information**


**Neuropilin 1 Rabbit pAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Neuropilin 1 Rabbit pAb - Images**






Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-NRP1/CD304 Polyclonal Antibody, Unconjugated(AP94626) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat small intestine tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-NRP1/CD304 Polyclonal Antibody, Unconjugated(AP94626) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Sample: Lane 1: Huvec (Human) Cell Lysate at 30 ug Lane 2: MDA-MB-231 (Human) Cell Lysate at 30 ug Lane 3: A549 (Human) Cell Lysate at 30 ug Lane 4: Hela (Human) Cell Lysate at 30 ug  
Primary: Anti-Neuropilin 1 (AP94626) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 103 kD Observed band size: 130 kD

### **Neuropilin 1 Rabbit pAb - Background**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.