

**NDUFS2 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP9769c****Specification**

---

**NDUFS2 Antibody (Center) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">O75306</a>
Other Accession	<a href="#">O641Y2</a> , <a href="#">O91WD5</a> , <a href="#">P17694</a>
Reactivity	<b>Human, Mouse</b>
Predicted	<b>Bovine, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>52546</b>
Antigen Region	<b>286-315</b>

**NDUFS2 Antibody (Center) - Additional Information****Gene ID** 4720**Other Names**

NADH dehydrogenase [ubiquinone] iron-sulfur protein 2, mitochondrial, Complex I-49kD, CI-49kD, NADH-ubiquinone oxidoreductase 49 kDa subunit, NDUFS2

**Target/Specificity**

This NDUFS2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 286-315 amino acids from the Central region of human NDUFS2.

**Dilution**WB~~1:1000  
IHC-P~~1:50~100**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NDUFS2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**NDUFS2 Antibody (Center) - Protein Information****Name** NDUFS2

**Function** Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (PubMed:[22036843](#), PubMed:[28031252](#), PubMed:[30922174](#)). Essential for the catalytic activity of complex I (PubMed:[22036843](#), PubMed:[30922174](#)). Essential for the assembly of complex I (By similarity). Redox-sensitive, critical component of the oxygen-sensing pathway in the pulmonary vasculature which plays a key role in acute pulmonary oxygen-sensing and hypoxic pulmonary vasoconstriction (PubMed:[30922174](#)). Plays an important role in carotid body sensing of hypoxia (By similarity). Essential for glia-like neural stem and progenitor cell proliferation, differentiation and subsequent oligodendrocyte or neuronal maturation (By similarity).

#### Cellular Location


Mitochondrion inner membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:Q641Y2}; Matrix side {ECO:0000250|UniProtKB:Q641Y2}

### NDUFS2 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### NDUFS2 Antibody (Center) - Images



Western blot analysis of lysates from A431, HeLa cell line, mouse brain and rat brain tissue lysate(from left to right), using NDUFS2 Antibody (Center)(Cat. #AP9769c). AP9769c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



NDUFS2 Antibody (Center) (Cat. #AP9769c) IHC analysis in formalin fixed and paraffin embedded hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the NDUFS2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **NDUFS2 Antibody (Center) - Background**

NDUFS2 is a core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (complex I). Mammalian mitochondrial complex I is composed of at least 43 different subunits, 7 of which are encoded by the mitochondrial genome, and the rest are the products of nuclear genes. The iron-sulfur protein fraction of complex I is made up of 7 subunits, including this gene product. Complex I catalyzes the NADH oxidation with concomitant ubiquinone reduction and proton ejection out of the mitochondria.

#### **NDUFS2 Antibody (Center) - References**

Saada, A., et al. *Am. J. Hum. Genet.* 84(6):718-727(2009) Wang, L., et al. *Cancer Epidemiol. Biomarkers Prev.* 17(12):3558-3566(2008) Starr, J.M., et al. *Mech. Ageing Dev.* 129(12):745-751(2008) Ban, M., et al. *PLoS ONE* 3 (8), E2891 (2008) Triepels, R.H., et al. *J. Biol. Chem.* 276(12):8892-8897(2001)