

ADCY4 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9875c

Specification

ADCY4 Antibody (Center) - Product Information

Application	WB, FC,E
Primary Accession	Q8NFM4
Other Accession	P26770 , Q91WF3
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	119794
Antigen Region	415-441

ADCY4 Antibody (Center) - Additional Information

Gene ID 196883

Other Names

Adenylate cyclase type 4, ATP pyrophosphate-lyase 4, Adenylate cyclase type IV, Adenylyl cyclase 4, ADCY4

Target/Specificity

This ADCY4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 415-441 amino acids from the Central region of human ADCY4.

Dilution

WB~~1:1000
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ADCY4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ADCY4 Antibody (Center) - Protein Information

Name ADCY4

Function Catalyzes the formation of the signaling molecule cAMP in response to G-protein signaling.

Cellular Location

Cell membrane; Multi-pass membrane protein. Cytoplasm

Tissue Location


Detected in the zona glomerulosa and the zona fasciculata in the adrenal gland (at protein level)

ADCY4 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADCY4 Antibody (Center) - Images



Western blot analysis of ADCY4 Antibody (Center) (Cat. #AP9875c) in HeLa cell line lysates (35ug/lane). ADCY4 (arrow) was detected using the purified Pab.



ADCY4 Antibody (Center) (Cat. #AP9875c) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ADCY4 Antibody (Center) - Background

This gene encodes a member of the family of adenylate cyclases, which are membrane-associated enzymes that catalyze the formation of the secondary messenger cyclic adenosine monophosphate (cAMP). Mouse studies show that adenylate cyclase 4, along with adenylate cyclases 2 and 3, is expressed in olfactory cilia, suggesting that several different adenylate cyclases may couple to olfactory receptors and that there may be multiple receptor-mediated mechanisms for the generation of cAMP signals.

ADCY4 Antibody (Center) - References

Rhim, J.H., et al. Aging Cell 5(6):451-461(2006)
 Jiang, G., et al. Am. J. Physiol. Endocrinol. Metab. 284 (4), E671-E678 (2003)
 Sunahara, R.K., et al. Mol. Interv. 2(3):168-184(2002)
 Ludwig, M.G., et al. J. Recept. Signal Transduct. Res. 22 (1-4), 79-110 (2002)
 Cote, M., et al. J. Clin. Endocrinol. Metab. 86(9):4495-4503(2001)