

Anti-TNFSF7 / CD27L / CD70 Reference Antibody (vorsetuzumab)

Recombinant Antibody Catalog # APR10097

Specification

Anti-TNFSF7 / CD27L / CD70 Reference Antibody (vorsetuzumab) - Product Information

Application FC, E, FTA
Primary Accession P32970
Reactivity Human
Clonality Monoclonal
Isotype IgG1
Calculated MW 146.1 KDa

Anti-TNFSF7 / CD27L / CD70 Reference Antibody (vorsetuzumab) - Additional Information

Target/Specificity
TNFSF7 / CD27L / CD70

Endotoxin

< 0.001EU/ µg,determined by LAL method.

Conjugation Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Anti-TNFSF7 / CD27L / CD70 Reference Antibody (vorsetuzumab) - Protein Information

Name CD70 (HGNC:11937)

Function

Expressed at the plasma membrane of B cells, it is the ligand of the CD27 receptor which is specifically expressed at the surface of T cells (PubMed:28011863, PubMed:28011864, PubMed:8387892). The CD70-CD27 signaling pathway mediates antigen-specific T cell activation and expansion which in turn provides immune surveillance of B cells (PubMed:28011863, PubMed:28011864).

Cellular Location

Cell membrane; Single-pass type II membrane protein




Anti-TNFSF7 / CD27L / CD70 Reference Antibody (vorsetuzumab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-TNFSF7 / CD27L / CD70 Reference Antibody (vorsetuzumab) - Images




Anti-TNFSF7 / CD27L / CD70 Reference Antibody (vorsetuzumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%




The purity of Anti-TNFSF7 / CD27L / CD70 Reference Antibody (vorsetuzumab)is more than 99.29% ,determined by SEC-HPLC.





HL-60 cells were stained with Anti-TNFSF7 / CD27L / CD70 Reference Antibody (vorsetuzumab) and negative control protein respectively, washed and then followed by PE and analyzed with FACS, EC150=0.06731 ug/mL



The endocytosis ratio vorsetuzumab by 786-O increased with the increase of antibody concentration, and the Internalization Rate (%) reached 55% at antibody concentration of 0.67 nM.