

# Anti-IL-3Ra / CD123 Reference Antibody (talacotuzumab-MMAE)

Recombinant Antibody Catalog # APR10128

# **Specification**

## Anti-IL-3Ra / CD123 Reference Antibody (talacotuzumab-MMAE) - Product Information

Application FC, E, FTA
Primary Accession P26951
Reactivity Human
Clonality Monoclonal
Isotype IgG1
Calculated MW 145 KDa

#### Anti-IL-3Ra / CD123 Reference Antibody (talacotuzumab-MMAE) - Additional Information

Target/Specificity IL-3Ra / CD123

**Endotoxin** 

< 0.001EU/ µg,determined by LAL method.

Conjugation

MMAE

**Expression system** 

CHO Cell

#### **Format**

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

### Anti-IL-3Ra / CD123 Reference Antibody (talacotuzumab-MMAE) - Protein Information

Name IL3RA (HGNC:6012)

Synonyms IL3R

#### **Function**

Cell surface receptor for IL3 expressed on hematopoietic progenitor cells, monocytes and B-lymphocytes that controls the production and differentiation of hematopoietic progenitor cells into lineage-restricted cells (PubMed:<a href="http://www.uniprot.org/citations/10527461" target="\_blank">10527461</a>). Ligand stimulation rapidly induces hetrodimerization with IL3RB, phosphorylation and enzyme activity of effector proteins such as JAK2 and PI3K that play a role in signaling cell proliferation and differentiation. Activation of JAK2 leads to STAT5-mediated transcriptional program (By similarity).

#### **Cellular Location**

Membrane; Single-pass type I membrane protein.




## Anti-IL-3Ra / CD123 Reference Antibody (talacotuzumab-MMAE) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

#### Anti-IL-3Ra / CD123 Reference Antibody (talacotuzumab-MMAE) - Images




Anti-IL-3Ra / CD123 Reference Antibody (talacotuzumab-MMAE) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-IL-3Ra / CD123 Reference Antibody (talacotuzumab-MMAE)is more than 95% ,determined by SEC-HPLC.





Immobilized human IL 3RA FC at 2  $~\mu g/mL$  can bind Anti-IL-3Ra / CD123 Reference Antibody (talacotuzumab-MMAE) []EC50=0.1647  $\mu g/mL$