

**Anti-CD38 Reference Antibody (isatuximab)
Recombinant Antibody
Catalog # APR10186****Specification**

Anti-CD38 Reference Antibody (isatuximab) - Product Information

Application	FC, E, FTA
Primary Accession	P28907
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145 KDa

Anti-CD38 Reference Antibody (isatuximab) - Additional Information**Target/Specificity**
CD38**Endotoxin**
< 0.001EU/ µg,determined by LAL method.**Conjugation**
Unconjugated**Expression system**
CHO Cell**Format**
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.**Anti-CD38 Reference Antibody (isatuximab) - Protein Information****Name** CD38**Function**
Synthesizes cyclic ADP-ribose (cADPR), a second messenger for glucose-induced insulin secretion (PubMed:7961800, PubMed:8253715). Synthesizes the Ca(2+) mobilizer nicotinate-adenine dinucleotide phosphate, NAADP(+), from 2'-phospho-cADPR and nicotinic acid, as well as from NADP(+) and nicotinic acid. At both pH 5.0 and pH 7.4 preferentially transforms 2'-phospho-cADPR into NAADP(+), while preferentially cleaving NADP(+) to cADPR and ADPRP rather than into NADDP(+) (PubMed:16690024). Has cADPR hydrolase activity (PubMed:7961800, PubMed:8253715).

Cellular Location

Cell surface. Membrane; Single-pass type II membrane protein


Tissue Location

Expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma.


Anti-CD38 Reference Antibody (isatuximab) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

Anti-CD38 Reference Antibody (isatuximab) - Images

Anti-CD38 Reference Antibody (isatuximab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-CD38 Reference Antibody (isatuximab) is more than 95% ,determined by SEC-HPLC.



Immobilized human CD38 Nhis at 2 µg/mL can bind Anti-CD38 Reference Antibody (isatuximab) \square EC50=0.006163 µg/mL