

Anti-OSMR Reference Antibody (vixarelimab)
Recombinant Antibody
Catalog # APR10201

Specification

Anti-OSMR Reference Antibody (vixarelimab) - Product Information

Application	FC, E, FTA
Primary Accession	Q99650
Reactivity	Cynomolgus, Human
Clonality	Monoclonal
Isotype	IgG4SP
Calculated MW	145 KDa

Anti-OSMR Reference Antibody (vixarelimab) - Additional Information

Target/Specificity
OSMR

Endotoxin
< 0.001EU/ µg,determined by LAL method.

Conjugation
Unconjugated

Expression system
CHO Cell

Format
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Anti-OSMR Reference Antibody (vixarelimab) - Protein Information

Name OSMR

Synonyms OSMRB

Function
Associates with IL31RA to form the IL31 receptor. Binds IL31 to activate STAT3 and possibly STAT1 and STAT5. Capable of transducing OSM-specific signaling events.

Cellular Location
Membrane; Single-pass type I membrane protein


Tissue Location
Expressed in keratinocytes (at protein level) (PubMed:21261663). Expressed at relatively high levels in all neural cells as well as fibroblast and epithelial cells (PubMed:8999038)

Anti-OSMR Reference Antibody (vixarelimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-OSMR Reference Antibody (vixarelimab) - Images




Anti-OSMR Reference Antibody (vixarelimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%




The purity of Anti-OSMR Reference Antibody (vixarelimab) is more than 96.9%, determined by SEC-HPLC.




Immobilized human OSMR His at 2 µg/mL can bind Anti-OSMR Reference Antibody (vixarelimab) □ EC50=0.0024 µg/mL



Human OSMR HEK293 cells were stained with Anti-OSMR Reference Antibody (vixarelimab) and negative control protein respectively, washed and then followed by PE and analyzed with FACS, EC261=0.0359 µg/mL



Anti-OSMR Reference Antibody (vixarelimab) P-STAT3 Assay was evaluated using NHDF. The IC50 was approximately 0.3493 nM.



Anti-OSMR Reference Antibody (vixarelimab) MCP-1 secretion was evaluated using NHDF. The IC50 was approximately 0.0812 nM.