

Anti-TGFb1 Reference Antibody (M7824)

Recombinant Antibody Catalog # APR10250

## Specification

# Anti-TGFb1 Reference Antibody (M7824) - Product Information

Application Primary Accession Reactivity Clonality Isotype Calculated MW FC, E, FTA <u>P01137</u> Cynomolgus, Human, Mouse Monoclonal IgG1 178 KDa

### Anti-TGFb1 Reference Antibody (M7824) - Additional Information

Target/Specificity TGFb1

**Endotoxin** < 0.001EU/ μg,determined by LAL method.

Conjugation Unconjugated

Expression system CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

### Anti-TGFb1 Reference Antibody (M7824) - Protein Information

Name TGFB1 (HGNC:11766)

Synonyms TGFB

Function

Transforming growth factor beta-1 proprotein: Precursor of the Latency-associated peptide (LAP) and Transforming growth factor beta-1 (TGF-beta-1) chains, which constitute the regulatory and active subunit of TGF-beta-1, respectively.

**Cellular Location** [Latency-associated peptide]: Secreted, extracellular space, extracellular matrix

**Tissue Location** 

Highly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498,




PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158)

### Anti-TGFb1 Reference Antibody (M7824) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

#### Anti-TGFb1 Reference Antibody (M7824) - Images




Anti-TGFb1 Reference Antibody (M7824) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90.7%



The purity of Anti-TGFb1 Reference Antibody (M7824)is more than 96.19% ,determined by SEC-HPLC.





Immobilized human PD L1 FC at 2  $\mu$ g/mL can bind Anti-TGFb1 Reference Antibody (M7824)]EC50=0.2227  $\mu$ g/mL