

**Anti-ICOS / CD278 Reference Antibody (vopratelimab)  
Recombinant Antibody  
Catalog # APR10287**

**Specification**

---

**Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Product Information**

Application	FC, E, FTA
Primary Accession	<a href="#">O9Y6W8</a>
Reactivity	Cynomolgus, Human, Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145.8 KDa

**Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Additional Information**

**Target/Specificity**  
ICOS / CD278

**Endotoxin**  
< 0.001EU/ µg,determined by LAL method.

**Conjugation**  
Unconjugated

**Expression system**  
CHO Cell

**Format**  
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

**Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Protein Information**

**Name** ICOS

**Synonyms** AILIM

**Function**  
Enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells. Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Does not up-regulate the production of interleukin- 2, but superinduces the synthesis of interleukin-10. Prevents the apoptosis of pre-activated T-cells. Plays a critical role in CD40- mediated class switching of immunoglobulin isotypes (By similarity).

**Cellular Location**  
[Isoform 1]: Cell membrane; Single- pass type I membrane protein

### Tissue Location


Activated T-cells. Highly expressed on tonsillar T- cells, which are closely associated with B-cells in the apical light zone of germinal centers, the site of terminal B-cell maturation Expressed at lower levels in thymus, lung, lymph node and peripheral blood leukocytes. Expressed in the medulla of fetal and newborn thymus

### Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-ICOS / CD278 Reference Antibody (vopratelimab) - Images



Anti-ICOS / CD278 Reference Antibody (vopratelimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-ICOS / CD278 Reference Antibody (vopratelimab) is more than 99.09%, determined by SEC-HPLC.