

**Anti-PDCD1 / PD-1 / CD279 Reference Antibody (finotonlimab)  
Recombinant Antibody  
Catalog # APR10334**

**Specification**

---

**Anti-PDCD1 / PD-1 / CD279 Reference Antibody (finotonlimab) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | FC, E, FTA             |
| Primary Accession | <a href="#">Q15116</a> |
| Reactivity        | Human                  |
| Clonality         | Monoclonal             |
| Isotype           | IgG4SP                 |
| Calculated MW     | 145 KDa                |

**Anti-PDCD1 / PD-1 / CD279 Reference Antibody (finotonlimab) - Additional Information**

**Target/Specificity**  
PDCD1 / PD-1 / CD279

**Endotoxin**  
< 0.001EU/ µg,determined by LAL method.

**Conjugation**  
Unconjugated

**Expression system**  
CHO Cell

**Format**  
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

**Storage**  
-80°C for 2 years under sterile conditions □ -20°C for 1 year under sterile conditions □ Avoid repeated freeze-thaw cycles.

**Anti-PDCD1 / PD-1 / CD279 Reference Antibody (finotonlimab) - Protein Information**

**Name** PDCD1 {ECO:0000303|PubMed:7851902, ECO:0000312|HGNC:HGNC:8760}

**Function**  
Inhibitory receptor on antigen activated T-cells that plays a critical role in induction and maintenance of immune tolerance to self (PubMed:<a href="http://www.uniprot.org/citations/21276005" target="\_blank">21276005</a>). Delivers inhibitory signals upon binding to ligands CD274/PDCD1L1 and CD273/PDCD1LG2 (PubMed:<a href="http://www.uniprot.org/citations/21276005" target="\_blank">21276005</a>). Following T-cell receptor (TCR) engagement, PDCD1 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). Suppresses T-cell activation through the recruitment of PTPN11/SHP-2: following ligand-binding, PDCD1 is phosphorylated within the ITSM

motif, leading to the recruitment of the protein tyrosine phosphatase PTPN11/SHP-2 that mediates dephosphorylation of key TCR proximal signaling molecules, such as ZAP70, PRKCQ/PKtheta and CD247/CD3zeta (By similarity).

#### Cellular Location


Cell membrane; Single-pass type I membrane protein

#### Anti-PDCD1 / PD-1 / CD279 Reference Antibody (finotonlimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-PDCD1 / PD-1 / CD279 Reference Antibody (finotonlimab) - Images



Anti-PDCD1 / PD-1 / CD279 Reference Antibody (finotonlimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-PDCD1 / PD-1 / CD279 Reference Antibody (finotonlimab) is more than 95%, determined by SEC-HPLC.