

**Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab)
Recombinant Antibody
Catalog # APR10343**

Specification

Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab) - Product Information

Application	FC, E, FTA
Primary Accession	P29965
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145.68 KDa

Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab) - Additional Information

Target/Specificity
TNFSF5 / CD40L / CD154

Endotoxin
< 0.001EU/ µg,determined by LAL method.

Conjugation
Unconjugated

Expression system
CHO Cell

Format
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab) - Protein Information

Name CD40LG

Synonyms CD40L, TNFSF5, TRAP

Function
Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed:[1280226](http://www.uniprot.org/citations/1280226), PubMed:[31331973](http://www.uniprot.org/citations/31331973)). Costimulates T-cell proliferation and cytokine production (PubMed:[8617933](http://www.uniprot.org/citations/8617933)). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed:[8617933](http://www.uniprot.org/citations/8617933)). Induces the activation of NF-kappa-B (PubMed:[15067037](http://www.uniprot.org/citations/15067037), PubMed:[31331973](http://www.uniprot.org/citations/31331973)

target="_blank">31331973). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed:15067037). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed:15067037). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity).

Cellular Location

Cell membrane; Single-pass type II membrane protein. Cell surface

Tissue Location


Specifically expressed on activated CD4+ T- lymphocytes

Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab) - Images



Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab) is more than 99% ,determined by SEC-HPLC.