

Anti-ICOS / CD278 Reference Antibody (MEDI-570)
Recombinant Antibody
Catalog # APR10439

Specification

Anti-ICOS / CD278 Reference Antibody (MEDI-570) - Product Information

Application	FC, E, FTA
Primary Accession	O9Y6W8
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145 KDa

Anti-ICOS / CD278 Reference Antibody (MEDI-570) - Additional Information

Target/Specificity
ICOS / CD278

Endotoxin
< 0.001EU/ µg,determined by LAL method.

Conjugation
Unconjugated

Expression system
CHO Cell

Format
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Anti-ICOS / CD278 Reference Antibody (MEDI-570) - Protein Information

Name ICOS

Synonyms AILIM

Function
Enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells. Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Does not up-regulate the production of interleukin- 2, but superinduces the synthesis of interleukin-10. Prevents the apoptosis of pre-activated T-cells. Plays a critical role in CD40- mediated class switching of immunoglobulin isotypes (By similarity).

Cellular Location
[Isoform 1]: Cell membrane; Single- pass type I membrane protein

Tissue Location


Activated T-cells. Highly expressed on tonsillar T- cells, which are closely associated with B-cells in the apical light zone of germinal centers, the site of terminal B-cell maturation Expressed at lower levels in thymus, lung, lymph node and peripheral blood leukocytes. Expressed in the medulla of fetal and newborn thymus

Anti-ICOS / CD278 Reference Antibody (MEDI-570) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

Anti-ICOS / CD278 Reference Antibody (MEDI-570) - Images



Anti-ICOS / CD278 Reference Antibody (MEDI-570) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-ICOS / CD278 Reference Antibody (MEDI-570) is more than 95% ,determined by SEC-HPLC.