

**Anti-CD28L / CD80 Reference Antibody (galiximab)  
Recombinant Antibody  
Catalog # APR10515****Specification**

---

**Anti-CD28L / CD80 Reference Antibody (galiximab) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | FC, E, FTA             |
| Primary Accession | <a href="#">P33681</a> |
| Reactivity        | Cynomolgus, Human      |
| Clonality         | Monoclonal             |
| Isotype           | IgG1                   |
| Calculated MW     | 145.24 KDa             |

**Anti-CD28L / CD80 Reference Antibody (galiximab) - Additional Information****Target/Specificity**  
CD28L / CD80**Endotoxin**  
< 0.001EU/ µg, determined by LAL method.**Conjugation**  
Unconjugated**Expression system**  
CHO Cell**Format**  
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.**Anti-CD28L / CD80 Reference Antibody (galiximab) - Protein Information****Name** CD80**Synonyms** CD28LG, CD28LG1, LAB7**Function**

Costimulatory molecule that belongs to the immunoglobulin superfamily that plays an important role in T-lymphocyte activation (PubMed: [38467718](http://www.uniprot.org/citations/38467718)). Acts as the primary auxiliary signal augmenting the MHC/TCR signal in naive T-cells together with the CD28 receptor which is constitutively expressed on the cell surface of T-cells (PubMed: [12196291](http://www.uniprot.org/citations/12196291)). In turn, activates different signaling pathways such as NF-kappa-B or MAPK leading to the production of different cytokines (PubMed: [10438913](http://www.uniprot.org/citations/10438913)). In addition, CD28/CD80 costimulatory signal stimulates glucose metabolism and ATP synthesis of T-cells by activating the PI3K/Akt signaling pathway (PubMed: <

href="http://www.uniprot.org/citations/12121659" target="\_blank">12121659</a>). Acts also as a regulator of PDL1/PDCD1 interactions to limit excess engagement of PDL1 and its inhibitory role in immune responses (PubMed:<a href="http://www.uniprot.org/citations/36727298" target="\_blank">36727298</a>). Expressed on B-cells, plays a critical role in regulating interactions between B-cells and T-cells in both early and late germinal center responses, which are crucial for the generation of effective humoral immune responses (By similarity).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein

#### Tissue Location


Expressed on activated B-cells, macrophages and dendritic cells

### Anti-CD28L / CD80 Reference Antibody (galiximab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

### Anti-CD28L / CD80 Reference Antibody (galiximab) - Images



Anti-CD28L / CD80 Reference Antibody (galiximab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-CD28L / CD80 Reference Antibody (galiximab) is more than 95.08% ,determined by SEC-HPLC.