

**Anti-IL-3Ra / CD123 Reference Antibody (SNG-CD123A)
Recombinant Antibody
Catalog # APR10542****Specification**

Anti-IL-3Ra / CD123 Reference Antibody (SNG-CD123A) - Product Information

Application	FC, E, FTA
Primary Accession	P26951
Reactivity	Cynomolgus, Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	146.14 KDa

Anti-IL-3Ra / CD123 Reference Antibody (SNG-CD123A) - Additional Information**Target/Specificity**
IL-3Ra / CD123**Endotoxin**
< 0.001EU/ µg,determined by LAL method.**Conjugation**
Unconjugated**Expression system**
CHO Cell**Format**
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.**Storage**
-80°C for 2 years under sterile conditions □ -20°C for 1 year under sterile conditions □ Avoid repeated freeze-thaw cycles.**Anti-IL-3Ra / CD123 Reference Antibody (SNG-CD123A) - Protein Information****Name** IL3RA ([HGNC:6012](#))**Synonyms** IL3R**Function**
Cell surface receptor for IL3 expressed on hematopoietic progenitor cells, monocytes and B-lymphocytes that controls the production and differentiation of hematopoietic progenitor cells into lineage-restricted cells (PubMed:10527461). Ligand stimulation rapidly induces heterodimerization with IL3RB, phosphorylation and enzyme activity of effector proteins such as JAK2 and PI3K that play a role in signaling cell proliferation and differentiation. Activation of JAK2 leads to STAT5-mediated

transcriptional program (By similarity).

Cellular Location


Membrane; Single-pass type I membrane protein.

Anti-IL-3Ra / CD123 Reference Antibody (SNG-CD123A) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IL-3Ra / CD123 Reference Antibody (SNG-CD123A) - Images



Anti-IL-3Ra / CD123 Reference Antibody (SNG-CD123A) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-IL-3Ra / CD123 Reference Antibody (SNG-CD123A) is more than 95.78%, determined by SEC-HPLC.