

**Anti-IL-5 Reference Antibody (depemokimab)  
Recombinant Antibody  
Catalog # APR10546****Specification**

---

**Anti-IL-5 Reference Antibody (depemokimab) - Product Information**

Application	FC, E, FTA
Primary Accession	<a href="#">P05113</a>
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145.8 KDa

**Anti-IL-5 Reference Antibody (depemokimab) - Additional Information****Target/Specificity**

IL-5

**Endotoxin**

&lt; 0.001EU/ µg,determined by LAL method.

**Conjugation**

Unconjugated

**Expression system**

CHO Cell

**Format**

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

**Anti-IL-5 Reference Antibody (depemokimab) - Protein Information****Name** IL5**Function**

Homodimeric cytokine expressed predominantly by T-lymphocytes and NK cells that plays an important role in the survival, differentiation, and chemotaxis of eosinophils (PubMed:<a href="http://www.uniprot.org/citations/2653458" target="\_blank">2653458</a>, PubMed:<a href="http://www.uniprot.org/citations/9010276" target="\_blank">9010276</a>). Acts also on activated and resting B-cells to induce immunoglobulin production, growth, and differentiation (By similarity). Mechanistically, exerts its biological effects through a receptor composed of IL5RA subunit and the cytokine receptor common subunit beta/CSF2RB (PubMed:<a href="http://www.uniprot.org/citations/1495999" target="\_blank">1495999</a>, PubMed:<a href="http://www.uniprot.org/citations/22528658" target="\_blank">22528658</a>). Binding to the receptor leads to activation of various kinases including LYN, SYK and JAK2 and thereby propagates signals through the RAS-MAPK and JAK-STAT5 pathways respectively (PubMed:<a href="http://www.uniprot.org/citations/7613138" target="\_blank">7613138</a>).

**Cellular Location**  
Secreted.


**Tissue Location**  
Present in peripheral blood mononuclear cells.

### Anti-IL-5 Reference Antibody (depemokimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-IL-5 Reference Antibody (depemokimab) - Images



Anti-IL-5 Reference Antibody (depemokimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-IL-5 Reference Antibody (depemokimab) is more than 98.75%, determined by SEC-HPLC.