

Anti-IL-9 Reference Antibody (enokizumab)

Recombinant Antibody Catalog # APR10578

Specification

Anti-IL-9 Reference Antibody (enokizumab) - Product Information

Application Primary Accession Reactivity Clonality Isotype Calculated MW FC, E, FTA P15248 Human Monoclonal IgG1 145 KDa

Anti-IL-9 Reference Antibody (enokizumab) - Additional Information

Target/Specificity IL-9

Endotoxin < 0.001EU/ μg,determined by LAL method.

Conjugation Unconjugated

Expression system CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Anti-IL-9 Reference Antibody (enokizumab) - Protein Information

Name IL9

Function

Multifunctional cytokine secreted mainly by T-helper 2 lymphocytes and also mast cells or NKT cells that plays important roles in the immune response against parasites (PubMed:29742432). Affects intestinal epithelial permeability and adaptive immunity (PubMed:29742432). In addition, induces the differentiation of specific T-cell subsets such as IL-17 producing helper T-cells (TH17) and also proliferation and differentiation of mast cells. Mechanistically, exerts its biological effects through a receptor composed of IL9R subunit and a signal transducing subunit IL2RG. Receptor stimulation results in the rapid activation of JAK1 and JAK3 kinase activities leading to STAT1, STAT3 and STAT5-mediated transcriptional programs. Induction of differentiation genes seems to be mediated by STAT1 alone, while protection of cells from apoptosis depends on STAT3 and STAT5.




Cellular Location Secreted.

Anti-IL-9 Reference Antibody (enokizumab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-IL-9 Reference Antibody (enokizumab) - Images



Anti-IL-9 Reference Antibody (enokizumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-IL-9 Reference Antibody (enokizumab)is more than 95% ,determined by SEC-HPLC.