

**Anti-F3 / Factor III / Tissue Factor / CD142 Reference Antibody (TNX-832)
Recombinant Antibody
Catalog # APR10714**

Specification

Anti-F3 / Factor III / Tissue Factor / CD142 Reference Antibody (TNX-832) - Product Information

Application	FC, E, FTA
Primary Accession	P13726
Reactivity	Cynomolgus, Human
Clonality	Monoclonal
Isotype	IgG4SP
Calculated MW	150 KDa

Anti-F3 / Factor III / Tissue Factor / CD142 Reference Antibody (TNX-832) - Additional Information

Target/Specificity

F3 / Factor III / Tissue Factor / CD142

Endotoxin

< 0.001EU/ µg,determined by LAL method.

Conjugation

Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Storage

-80°C for 2 years under sterile conditions □ -20°C for 1 year under sterile conditions □ Avoid repeated freeze-thaw cycles.

Anti-F3 / Factor III / Tissue Factor / CD142 Reference Antibody (TNX-832) - Protein Information

Name F3

Function

Initiates blood coagulation by forming a complex with circulating factor VII or VIIa. The [TF:VIIa] complex activates factors IX or X by specific limited proteolysis. TF plays a role in normal hemostasis by initiating the cell-surface assembly and propagation of the coagulation protease cascade.

Cellular Location

[Isoform 1]: Membrane; Single-pass type I membrane protein


Tissue Location

Lung, placenta and pancreas.


Anti-F3 / Factor III / Tissue Factor / CD142 Reference Antibody (TNX-832) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-F3 / Factor III / Tissue Factor / CD142 Reference Antibody (TNX-832) - Images

Anti-F3 / Factor III / Tissue Factor / CD142 Reference Antibody (TNX-832) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-F3 / Factor III / Tissue Factor / CD142 Reference Antibody (TNX-832) is more than 95% ,determined by SEC-HPLC.