

**Anti-IL-18 Reference Antibody (ABT-325)
Recombinant Antibody
Catalog # APR10737****Specification**

Anti-IL-18 Reference Antibody (ABT-325) - Product Information

| | |
|-------------------|------------------------|
| Application | FC, E, FTA |
| Primary Accession | Q14116 |
| Reactivity | Human |
| Clonality | Monoclonal |
| Isotype | IgG1 |
| Calculated MW | 146.28 KDa |

Anti-IL-18 Reference Antibody (ABT-325) - Additional Information**Target/Specificity**
IL-18**Endotoxin**
< 0.001EU/ µg, determined by LAL method.**Conjugation**
Unconjugated**Expression system**
CHO Cell**Format**
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.**Anti-IL-18 Reference Antibody (ABT-325) - Protein Information****Name** IL18 ([HGNC:5986](#))**Synonyms** IGIF, IL1F4**Function**
Pro-inflammatory cytokine primarily involved in epithelial barrier repair, polarized T-helper 1 (Th1) cell and natural killer (NK) cell immune responses (PubMed: [10653850](http://www.uniprot.org/citations/10653850)). Upon binding to IL18R1 and IL18RAP, forms a signaling ternary complex which activates NF-kappa-B, triggering synthesis of inflammatory mediators (PubMed: [14528293](http://www.uniprot.org/citations/14528293), PubMed: [25500532](http://www.uniprot.org/citations/25500532), PubMed: [37993714](http://www.uniprot.org/citations/37993714)). Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells and natural killer (NK) cells (PubMed: [10653850](http://www.uniprot.org/citations/10653850))

target="_blank">10653850). Involved in transduction of inflammation downstream of pyroptosis: its mature form is specifically released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744).

Cellular Location

Cytoplasm, cytosol. Secreted. Note=The precursor is cytosolic (PubMed:33883744). In response to inflammasome-activating signals, cleaved and secreted (PubMed:33883744, PubMed:37993712, PubMed:37993714). Mature form is secreted and released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744, PubMed:37993714). In contrast, the precursor form is not released, due to the presence of an acidic region that is proteolytically removed by CASP1, CASP4 or CASP5 during maturation (PubMed:33883744, PubMed:37993714). The secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10 (PubMed:32272059).

Tissue Location


[Isoform 2]: Expressed in ovarian carcinoma but undetectable in normal ovarian epithelial cells. Resistant to proteolytic activation by caspase-1 and -4

Anti-IL-18 Reference Antibody (ABT-325) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IL-18 Reference Antibody (ABT-325) - Images



Anti-IL-18 Reference Antibody (ABT-325) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-IL-18 Reference Antibody (ABT-325) is more than 95% ,determined by SEC-HPLC.