

Anti-TIGIT Reference Antibody (domvanalimab)

Recombinant Antibody Catalog # APR10741

Specification

Anti-TIGIT Reference Antibody (domvanalimab) - Product Information

Application FC, E, FTA Primary Accession Q495A1

Reactivity Cynomolgus, Human

Clonality Monoclonal

Isotype IgG1

Calculated MW 145.12 KDa

Anti-TIGIT Reference Antibody (domvanalimab) - Additional Information

Target/Specificity

TIGIT

Endotoxin

 $< 0.001EU/ \mu g$, determined by LAL method.

Conjugation Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Anti-TIGIT Reference Antibody (domvanalimab) - Protein Information

Name TIGIT

Synonyms VSIG9, VSTM3

Function

Binds with high affinity to the poliovirus receptor (PVR) which causes increased secretion of IL10 and decreased secretion of IL12B and suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed at low levels on peripheral memory and regulatory CD4+ T-cells and NK cells and is up-regulated following activation of these cells (at protein level)




Anti-TIGIT Reference Antibody (domvanalimab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-TIGIT Reference Antibody (domvanalimab) - Images



Anti-TIGIT Reference Antibody (domvanalimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-TIGIT Reference Antibody (domvanalimab)is more than 99.65% ,determined by SEC-HPLC.