

**Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151)
Recombinant Antibody
Catalog # APR10799**

Specification

Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151) - Product Information

Application	FC, E, FTA
Primary Accession	P48509
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145 KDa

Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151) - Additional Information

Target/Specificity

CD151

Endotoxin

< 0.001EU/ µg,determined by LAL method.

Conjugation

Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151) - Protein Information

Name CD151

Synonyms TSPAN24

Function

Structural component of specialized membrane microdomains known as tetraspanin-enriched microdomains (TERMs), which act as platforms for receptor clustering and signaling. Plays a role in various cellular and molecular mechanism through its association with both integrin and non-integrin proteins. These interactions facilitate critical cellular functions, including cell-to-cell communication, wound healing, platelet aggregation, trafficking, cell motility, and angiogenesis (PubMed:17045834, PubMed:24723389, PubMed:31488507). Via interaction with JAM-A/F11R and integrin ITGA3:ITGB1, promotes the recruitment of signaling

molecules such as RAC1, CDC42 and RhoGTPases to facilitate the polarization of epithelial cells and the reorganization of the actin cytoskeleton, which are critical steps in cell migration process (PubMed:22843693, PubMed:35067832). Regulates the glycosylation pattern of ITGA3:ITGB1 thereby modulating its activity (PubMed:18852263). Plays an essential role in the maintenance of central laminin-binding integrin ITGA6:ITGB4-containing adhesion complexes (PubMed:31488507). Essential for the proper assembly of the glomerular and tubular basement membranes in kidney (PubMed:15265795). Contributes to T-cell activation by modulating integrin signaling leading to activation of downstream targets PTK2 and MAPK1/MAPK3 (PubMed:24723389).

Cellular Location

Cell membrane; Multi-pass membrane protein Note=Relocalizes to the immune synapse in T-cells upon activation

Tissue Location


Expressed in a variety of tissues including vascular endothelium and epidermis. Expressed on erythroid cells, with a higher level of expression in erythroid precursors than on mature erythrocytes (PubMed:15265795). Acts as a sensitive T-cell activation marker (PubMed:32978478).

Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151) - Images



Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151) is more than 95% ,determined by SEC-HPLC.