

**Anti-c-RET Reference Antibody (Regeneron patent anti-RET)  
Recombinant Antibody  
Catalog # APR10848****Specification**

---

**Anti-c-RET Reference Antibody (Regeneron patent anti-RET) - Product Information**

Application	FC, E, FTA
Primary Accession	<a href="#">P07949</a>
Reactivity	Cynomolgus, Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145 KDa

**Anti-c-RET Reference Antibody (Regeneron patent anti-RET) - Additional Information****Target/Specificity**

c-RET

**Endotoxin**

&lt; 0.001EU/ µg,determined by LAL method.

**Conjugation**

Unconjugated

**Expression system**

CHO Cell

**Format**

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

**Anti-c-RET Reference Antibody (Regeneron patent anti-RET) - Protein Information****Name** RET {ECO:0000303|PubMed:2660074, ECO:0000312|HGNC:HGNC:9967}**Function**

Receptor tyrosine-protein kinase involved in numerous cellular mechanisms including cell proliferation, neuronal navigation, cell migration, and cell differentiation in response to glia cell line- derived growth family factors (GDNF, NRTN, ARTN, PSPN and GDF15) (PubMed:<a href="http://www.uniprot.org/citations/20064382" target="\_blank">20064382</a>, PubMed:<a href="http://www.uniprot.org/citations/20616503" target="\_blank">20616503</a>, PubMed:<a href="http://www.uniprot.org/citations/20702524" target="\_blank">20702524</a>, PubMed:<a href="http://www.uniprot.org/citations/21357690" target="\_blank">21357690</a>, PubMed:<a href="http://www.uniprot.org/citations/21454698" target="\_blank">21454698</a>, PubMed:<a href="http://www.uniprot.org/citations/24560924" target="\_blank">24560924</a>, PubMed:<a href="http://www.uniprot.org/citations/28846097" target="\_blank">28846097</a>, PubMed:<a href="http://www.uniprot.org/citations/28846099" target="\_blank">28846099</a>, PubMed:<a href="http://www.uniprot.org/citations/28953886" target="\_blank">28953886</a>, PubMed:<a

href="http://www.uniprot.org/citations/31118272" target="\_blank">31118272</a>). In contrast to most receptor tyrosine kinases, RET requires not only its cognate ligands but also coreceptors, for activation (PubMed:<a href="http://www.uniprot.org/citations/21994944" target="\_blank">21994944</a>, PubMed:<a href="http://www.uniprot.org/citations/23333276" target="\_blank">23333276</a>, PubMed:<a href="http://www.uniprot.org/citations/28846097" target="\_blank">28846097</a>, PubMed:<a href="http://www.uniprot.org/citations/28846099" target="\_blank">28846099</a>, PubMed:<a href="http://www.uniprot.org/citations/28953886" target="\_blank">28953886</a>). GDNF ligands (GDNF, NRTN, ARTN, PSPN and GDF15) first bind their corresponding GDNFR coreceptors (GFRA1, GFRA2, GFRA3, GFRA4 and GFRAL, respectively), triggering RET autophosphorylation and activation, leading to activation of downstream signaling pathways, including the MAPK- and AKT-signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/21994944" target="\_blank">21994944</a>, PubMed:<a href="http://www.uniprot.org/citations/23333276" target="\_blank">23333276</a>, PubMed:<a href="http://www.uniprot.org/citations/24560924" target="\_blank">24560924</a>, PubMed:<a href="http://www.uniprot.org/citations/25242331" target="\_blank">25242331</a>, PubMed:<a href="http://www.uniprot.org/citations/28846097" target="\_blank">28846097</a>, PubMed:<a href="http://www.uniprot.org/citations/28846099" target="\_blank">28846099</a>, PubMed:<a href="http://www.uniprot.org/citations/28953886" target="\_blank">28953886</a>). Acts as a dependence receptor via the GDNF-GFRA1 signaling: in the presence of the ligand GDNF in somatotrophs within pituitary, promotes survival and down regulates growth hormone (GH) production, but triggers apoptosis in absence of GDNF (PubMed:<a href="http://www.uniprot.org/citations/20616503" target="\_blank">20616503</a>, PubMed:<a href="http://www.uniprot.org/citations/21994944" target="\_blank">21994944</a>). Required for the molecular mechanisms orchestration during intestine organogenesis via the ARTN-GFRA3 signaling: involved in the development of enteric nervous system and renal organogenesis during embryonic life, and promotes the formation of Peyer's patch-like structures, a major component of the gut-associated lymphoid tissue (By similarity). Mediates, through interaction with GDF15-receptor GFRAL, GDF15-induced cell-signaling in the brainstem which triggers an aversive response, characterized by nausea, vomiting, and/or loss of appetite in response to various stresses (PubMed:<a href="http://www.uniprot.org/citations/28846097" target="\_blank">28846097</a>, PubMed:<a href="http://www.uniprot.org/citations/28846099" target="\_blank">28846099</a>, PubMed:<a href="http://www.uniprot.org/citations/28953886" target="\_blank">28953886</a>). Modulates cell adhesion via its cleavage by caspase in sympathetic neurons and mediates cell migration in an integrin (e.g. ITGB1 and ITGB3)-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/20702524" target="\_blank">20702524</a>, PubMed:<a href="http://www.uniprot.org/citations/21357690" target="\_blank">21357690</a>). Also active in the absence of ligand, triggering apoptosis through a mechanism that requires receptor intracellular caspase cleavage (PubMed:<a href="http://www.uniprot.org/citations/21357690" target="\_blank">21357690</a>). Triggers the differentiation of rapidly adapting (RA) mechanoreceptors (PubMed:<a href="http://www.uniprot.org/citations/20064382" target="\_blank">20064382</a>). Involved in the development of the neural crest (By similarity). Regulates nociceptor survival and size (By similarity). Phosphorylates PTK2/FAK1 (PubMed:<a href="http://www.uniprot.org/citations/21454698" target="\_blank">21454698</a>).

### Cellular Location

Cell membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein Note=Predominantly located on the plasma membrane (PubMed:23333276, PubMed:9575150). In the presence of SORL1 and GFRA1, directed to endosomes (PubMed:23333276).


### Anti-c-RET Reference Antibody (Regeneron patent anti-RET) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-c-RET Reference Antibody (Regeneron patent anti-RET) - Images



Anti-c-RET Reference Antibody (Regeneron patent anti-RET) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-c-RET Reference Antibody (Regeneron patent anti-RET) is more than 95%, determined by SEC-HPLC.