

**Anti-MMP9 Reference Antibody (Yeda patent anti-MMP-9)  
Recombinant Antibody  
Catalog # APR10979****Specification**

---

**Anti-MMP9 Reference Antibody (Yeda patent anti-MMP-9) - Product Information**

Application	FC, E, FTA
Primary Accession	<a href="#">P14780</a>
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	IgG2SA
Calculated MW	144.76 KDa

**Anti-MMP9 Reference Antibody (Yeda patent anti-MMP-9) - Additional Information****Target/Specificity**  
MMP9**Endotoxin**

< 0.001EU/ µg,determined by LAL method.

**Conjugation**

Unconjugated

**Expression system**

CHO Cell

**Format**

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

**Anti-MMP9 Reference Antibody (Yeda patent anti-MMP-9) - Protein Information****Name** MMP9**Synonyms** CLG4B**Function**

Matrix metalloproteinase that plays an essential role in local proteolysis of the extracellular matrix and in leukocyte migration (PubMed:[12879005](http://www.uniprot.org/citations/12879005), PubMed:[1480034](http://www.uniprot.org/citations/1480034), PubMed:[2551898](http://www.uniprot.org/citations/2551898)). Could play a role in bone osteoclastic resorption (By similarity). Cleaves KiSS1 at a Gly-I-Leu bond (PubMed:[12879005](http://www.uniprot.org/citations/12879005)). Cleaves NINJ1 to generate the Secreted ninjurin-1 form (PubMed:[32883094](http://www.uniprot.org/citations/32883094)). Cleaves type IV and type V collagen into large C-terminal three quarter fragments and shorter N-terminal one quarter fragments (PubMed:[1480034](http://www.uniprot.org/citations/1480034))

target="\_blank">1480034</a>). Degrades fibronectin but not laminin or Pz-peptide.

#### Cellular Location

Secreted, extracellular space, extracellular matrix

#### Tissue Location


Detected in neutrophils (at protein level) (PubMed:7683678). Produced by normal alveolar macrophages and granulocytes.

### Anti-MMP9 Reference Antibody (Yeda patent anti-MMP-9) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-MMP9 Reference Antibody (Yeda patent anti-MMP-9) - Images



Anti-MMP9 Reference Antibody (Yeda patent anti-MMP-9) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-MMP9 Reference Antibody (Yeda patent anti-MMP-9) is more than 95%, determined by SEC-HPLC.