

**Anti-PCSK9 Reference Antibody (Merck patent anti-PCSK9)  
Recombinant Antibody  
Catalog # APR11004**

**Specification**

---

**Anti-PCSK9 Reference Antibody (Merck patent anti-PCSK9) - Product Information**

Application	FC, E, FTA
Primary Accession	<a href="#">Q8NBP7</a>
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	150 KDa

**Anti-PCSK9 Reference Antibody (Merck patent anti-PCSK9) - Additional Information**

**Target/Specificity**  
PCSK9

**Endotoxin**  
< 0.001EU/ µg,determined by LAL method.

**Conjugation**  
Unconjugated

**Expression system**  
CHO Cell

**Format**  
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

**Anti-PCSK9 Reference Antibody (Merck patent anti-PCSK9) - Protein Information**

**Name** PCSK9

**Synonyms** NARC1

**Function**  
Crucial player in the regulation of plasma cholesterol homeostasis. Binds to low-density lipid receptor family members: low density lipoprotein receptor (LDLR), very low density lipoprotein receptor (VLDLR), apolipoprotein E receptor (LRP1/APOER) and apolipoprotein receptor 2 (LRP8/APOER2), and promotes their degradation in intracellular acidic compartments (PubMed:<a href="http://www.uniprot.org/citations/18039658" target="\_blank">18039658</a>). Acts via a non- proteolytic mechanism to enhance the degradation of the hepatic LDLR through a clathrin LDLRAP1/ARH-mediated pathway. May prevent the recycling of LDLR from endosomes to the cell surface or direct it to lysosomes for degradation. Can induce ubiquitination of LDLR leading to its subsequent degradation (PubMed:<a href="http://www.uniprot.org/citations/17461796" target="\_blank">17461796</a>, PubMed:<a href="http://www.uniprot.org/citations/18197702" target="\_blank">18197702</a>).

target="\_blank">18197702</a>, PubMed:<a href="http://www.uniprot.org/citations/18799458" target="\_blank">18799458</a>, PubMed:<a href="http://www.uniprot.org/citations/22074827" target="\_blank">22074827</a>). Inhibits intracellular degradation of APOB via the autophagosome/lysosome pathway in a LDLR-independent manner. Involved in the disposal of non-acetylated intermediates of BACE1 in the early secretory pathway (PubMed:<a href="http://www.uniprot.org/citations/18660751" target="\_blank">18660751</a>). Inhibits epithelial Na(+) channel (ENaC)-mediated Na(+) absorption by reducing ENaC surface expression primarily by increasing its proteasomal degradation. Regulates neuronal apoptosis via modulation of LRP8/APOER2 levels and related anti-apoptotic signaling pathways.

#### Cellular Location

Cytoplasm. Secreted. Endosome. Lysosome. Cell surface. Endoplasmic reticulum. Golgi apparatus. Note=Autocatalytic cleavage is required to transport it from the endoplasmic reticulum to the Golgi apparatus and for the secretion of the mature protein Localizes to the endoplasmic reticulum in the absence of LDLR and colocalizes to the cell surface and to the endosomes/lysosomes in the presence of LDLR. The sorting to the cell surface and endosomes is required in order to fully promote LDLR degradation

#### Tissue Location


Expressed in neuro-epithelioma, colon carcinoma, hepatic and pancreatic cell lines, and in Schwann cells

### Anti-PCSK9 Reference Antibody (Merck patent anti-PCSK9) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-PCSK9 Reference Antibody (Merck patent anti-PCSK9) - Images



Anti-PCSK9 Reference Antibody (Merck patent anti-PCSK9) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-PCSK9 Reference Antibody (Merck patent anti-PCSK9) is more than 95% ,determined by SEC-HPLC.