

DFF45 Antibody
Catalog # ASC10023**Specification**

DFF45 Antibody - Product Information

Application	WB
Primary Accession	O00273
Other Accession	NP_004392 , 1676
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	45 and 35 kDa KDa
Application Notes	DFF45 antibody can be used for detection of DFF45, DFF35, and one the cleaved fragment by Western blot at 1:1000 to 1:2000 dilution. 45 and 35 kDa bands can be detected in non-apoptotic cells.

DFF45 Antibody - Additional InformationGene ID **1676****Other Names**

DFF45 Antibody: DFF1, ICAD, DFF-45, DFF1, DFF45, H13, DNA fragmentation factor subunit alpha, DNA fragmentation factor 45 kDa subunit, DNA fragmentation factor, 45kDa, alpha polypeptide

Target/Specificity

DFF45 antibody was raised against a 20 amino acid peptide near the amino terminus of human DFF45.

The immunogen is located within the first 50 amino acids of DFF45.

Reconstitution & Storage

DFF45 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

DFF45 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

DFF45 Antibody - Protein Information**Name** DFFA**Synonyms** DFF1, DFF45**Function**

Inhibitor of the caspase-activated DNase (DFF40).

Cellular Location


Cytoplasm.

DFF45 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DFF45 Antibody - Images



Western blot analysis of DFF45 in HeLa cell lysate with DFF45 antibody at (A) 1 and (B) 2 µg/mL.

DFF45 Antibody - Background

DFF45 Antibody: Apoptosis is related to many diseases and induced by a family of cell death receptors and their ligands. Cell death signals are transduced by death domain containing adapter molecules and members of the caspase family of proteases. These death signals finally cause the degradation of chromosomal DNA by activated DNase. A human 45 kDa DNA fragmentation factor (DFF45) was identified recently that was cleaved by caspase-3 during apoptosis. Mouse homologue of human DFF45 was identified as a DNase inhibitor designated ICAD. DFF45/ICAD have short forms that were termed DFF35 and ICADs, respectively. Upon cleavage of DFF45/ICAD, the caspase activated deoxyribonuclease (DFF40/CAD) is released and activated and eventually causes the degradation of DNA in the nuclei. Therefore, the cleavage of DFF45/ICAD, which causes DFF40/CAD activation and DNA degradation, is the hallmark of apoptotic cell death.

DFF45 Antibody - References

Liu X, Zou H, Slaughter C, Wang X. DFF, a heterodimeric protein that functions downstream of caspase-3 to trigger DNA fragmentation during apoptosis. *Cell* 1997;89:175-184
Enari M, Sakahira H, Yokoyama H, Okawa K, Iwamatsu A, Nagata S. A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD. *Nature* 1998;391:43-50

Sakahira H, Enari M, Nagata S. Cleavage of CAD inhibitor in CAD activation and DNA degradation during apoptosis. *Nature* 1998;391:96-99

Gu J, Dong RP, Zhang C, McLaughlin DF, Wu MX, Schlossman SF. Functional interaction of DFF35 and DFF45 with caspase-activated DNA fragmentation nuclease DFF40. *J Biol Chem* 1999;274:20759-62